EMERALD COAST Obstetrics and Gynecology



PRENATAL BOOK

103 E. 23rd Street
Panama City, FL 32405
(850) 769-0338
www.emeraldcoastobgyn.com

Emerald Coast Providers



Henry Breland, MD FACOG



George Ramie, DO FACOOG



Samuel Wolf, DO FACOG



Toni Pennington, MD FACOG



Kimberly W. Bain, MD FACOG



Justo Maqueira, Jr., MD FACOG



Jacob Martin, MD FACOG



Heather Pate, MD Board Eligible



Michelle Lang MSN, APRN-C



MSN, APRN-C



Amanda Rodriguez MSN, APRN-C, CNM



Shawn Brooking DNP, APRN, CNM



Kim McConnell MSN, APRN, CNM



Shelrel Berry MSN, APRN, CNM



Cristina Stevenson MSN, APRN, CNM

PATIENT ID:	
-------------	--

I acknowledge that I received a copy of the



PRENATAL BOOK

PRINT NAME:		
Cian	Data	



Table of Contents



Congratulations/ Patient Portal	3
Telephone Directory	
Community Services	8
Providers and Appointments	10
Gulf Coast Medical Center	17
Laboratory Tests	19
Ultrasound	23
Approved Medications	
Prenatal Education	26
Morning Sickness	27
Warning Signs During Pregnancy	29
Nutrition During Pregnancy	
Smoking	34
Important Immunizations During Pregnancy	36
Influenza (flu) Infection and Pregnancy	37
Herpes Infections & Pregnancy	
Headache in Pregnancy	
Ligament Pain	
Back Pain During Pregnancy	43
Exercise During Pregnancy	46
Travel During Pregnancy	
Fetal Kick Counts	49
Urinary Tract Infections	50
Pre-term Labor	52
Blood Pressure Problems in Pregnancy	
Gestational Diabetes Mellitus	56
Abdominal Trauma	
Additional Fetal Surveillance	
Group B Strep	59
Vaginal Birth After Cesarean (VBAC)	60
Selecting a Pediatrician	
Induction of Labor	62
Broken Water	65
True Labor	
Labor Epidurals	
Hospital Packing List	
Circumcision	
Postpartum Information	74
Family Planning	
Interesting Websites	
Pregnancy Myths	
Core Blood Banking	
Notes	
Sponsor Information	89

This book belongs to:

Name:	 	
Phone:	 	
Due Date:		

EMERALD COAST OB/GYN OBSTETRIC PHILOSOPHY

As obstetric providers, we will always work diligently to improve the lives of the families we serve. However, we cannot accomplish the most optimal pregnancy outcome without the constant and careful cooperation of the most important member of our team — the expectant mother herself.

All rights reserved. Published by Emerald Coast OB/GYN, June, 2008 11th edition, October 2023 Editor: Samuel Wolf, DO



Congratulations!



Welcome to our obstetric services! Emerald Coast OB/GYN has a long tradition of providing the very best of care to expectant families living in Bay County and surrounding counties in northwest Florida. We are pleased that you have chosen us to help you throughout this important time in your life.

We have prepared this booklet to introduce our practice to you and to provide you with important information and helpful answers to many questions that will come up during your pregnancy.

Be sure to slip our little book into your bag and carry it with you at all times. You will find that you will often refer to the information contained in this book throughout your pregnancy.

This book may also be found on our website <u>www.emeraldcoastobgyn.com</u>. A complimentary copy will be given to you at your first visit. If you lose your book, you may acquire another book for \$8.

Register on the Patient Portal

Now you can update your patient information, check test results, pay your bill, request appointments, send and receive messages from our staff by accessing the Patient Portal. Please take time to activate your account.

First Time Registration: Go to the website www.emeraldcoastobgyn.com. At the top under "Patient Area" click on Patient Portal. On the right of the screen click the blue link that says "Sign Up Today".

- 1. Find Your Patient Record: Enter your name, birth date, email and phone number, Click "continue". Note: Enter the phone number on file from your first visit. A confirmation code will be presented on screen. You will need that number to complete the activation.
- **2. Verify Your Identity:** To ensure your privacy, we must verify your identity with a phone call, text, or email. Enter the phone number you would like to receive your call on and click "send code". When you receive the call, answer and follow the directions. Enter the confirmation code you were given on your call into the computer. When verification is complete you will continue the registration process on your computer screen.
- **3. Create Your Password:** You will be prompted to create a password for future logins on the patient portal. Please keep this password somewhere safe.

AFTER HOURS

We have a voicemail system that will pick up your call after hours and on the weekends. The system will direct your message to our on-call provider. If you ever call our office (850-769-0338) and no one returns your call, please keep in mind that your back-up option is Gulf Coast Hospital's Labor & Delivery Unit at (850-747-7700). Our providers are on call for urgent problems. If you are calling for a prescription refill or a non-urgent matter, please send a portal message on the patient portal or call back during normal business hours.

BUSINESS HOURS

Patient Portal: You can send secure messages to our clinical staff by logging into your patient portal account and go to "messages" and compose message".

Triage Line: OPTION # 5. We believe we have the busiest and best obstetrical call center in Northwest Florida. If you are unable to find the answer to your question in your book, please have it available for reference during your call. If you call our office during business hours and need advice, prescription refills, or want a question answered, your call will be first taken by a member of our staff. If all staff are busy assisting others, your call will automatically go to the voice mail system, and a member of our staff will be back in-touch with you as soon as possible and will be triaged appropriately.

Other Office Calls: If you are calling our office for any other reason, please listen carefully for prompts to all departments. If you do not hear an option that best fits your needs, please press the operator option and your call will be directed to the person who can best help you.

LOCATIONS

Our main office is located at 103 E. 23rd Street, Panama City, FL 32405. See our website for map and directions.

SCHEDULING

We cannot guarantee which provider will be present at the time of your delivery.

When you are checking out with our scheduling department, you are welcome to indicate which provider you would like to see for your next visit. If you require, or would like a labor induction when appropriate, you are welcome to request a specific provider for delivery, and we will try our best to accommodate your choice.

OFFICE VISITS

Prenatal appointments will generally be scheduled at monthly intervals up to 28 weeks gestation, then every two weeks from 28 – 36 weeks gestation, and then weekly until delivery. Should you have complications or become high risk, more frequent visits may become necessary.

CHECKING IN

Make sure you sign in at the front desk for all appointments. Please sign in 10-15 minutes prior to your appointment time so we can keep you on schedule. If you arrive late to your ultrasound visit you may be asked to reschedule to another time and date. If at any time during your pregnancy you have a change in insurance, benefits, or contact information, please let us know immediately.

URINE SPECIMEN

We check almost every patient's urine in our office. It is important that when you give a urine specimen that it is a "clean catch". In order to perform a clean catch, separate your labia and urinate for a second or two in the toilet. Introduce the urine cup in mid-stream without allowing the stream to touch the vagina or labia. This prevents false positive on your urine results.

FINANCIAL COUNSELOR/INSURANCE

We are very proud to be able to offer our patients excellent financial counseling services. We have a full-time financial counselor who is very knowledgeable about health insurance reimbursement or will be of valuable help in arranging a self-pay payment plan if you will benefit from one. She can also help to identify your payment options that can be a huge relief to your budget during this important period of your life. If at any time during your pregnancy you have a change in insurance, benefits, or contact information, please let us know PRIOR to your next scheduled appointment. Call to make an appointment with her to sit down and discuss your individual needs and concerns.

NICA

We are a NICA participating practice. The Florida Birth-Related Neurological Injury Compensation Association (NICA) was created by the Florida Legislature in 1988. NICA is a statutory organization that manages the Florida Birth Related Neurological Injury Compensation Plan used to pay for the care of infants born with certain neurological injuries. This plan is available to eligible families statewide without litigation. By eliminating costly legal proceedings, and through professional management of its disbursements, NICA ensures that birth-injured infants receive the care they need while reducing the financial burden on medical providers and families. There are restrictions to eligibility for NICA coverage.

FMLA/DISABILITY PAPERWORK

If you have FMLA paperwork that needs to be filled out for your maternity leave, please leave it with the front desk staff. Please allow 7-10 business days for this to be completed. We will call you when it is ready for pickup, or we are more than happy to fax it for you.

HEALTHY START SERVICES

We are happy to provide Healthy Start Services to our prenatal patients. These services include care coordination, smoking cessation, parenting and child development education, bereavement counseling, breastfeeding education, car seat safety education, stress managment courses and community referral. There is no cost to the family and this program is not based on income. You will fill out a Healthy Start risk assessment questionnaire to determine eligibility or you can request a specific service. The goals of Healthy Start are to improve pregnancy outcomes, to reduce infant mortality rates and to promote healthy growth and development. For more information please see Jennifer Bunnell when you are in our office or call her at 850-769-0338 ext. 180.

NATURAL DISASTERS

If the office should cose due to a natural disaster (such as a hurricane), our currently scheduled patients will be contacted about rescheduling. Also, we will post important updates and office closures on our Facebook and Instagram.

IMPORTANT UPDATES

To keep up to date with the latest information regarding pregnancy care, like us on social media pages.

We will periodically send out updates on important issues that may come up related to pregnancy.

www.facebook.com/emeraldcoastobgyn

Telephone Directory



Bay County Health Department 597 W. 11 th St. Panama City, FL	850-872-4720
Bay County Healthy Start and Lactation Support 2101 Northside Dr. Unit #101, Panama City FL 32405	850-872-4130
Domestic Violence Hotline	1-800-500-1119 Florida Relay 711
Emerald Coast Obstetrics and Gynecology	850-769-0338
Emergency Police / Fire / Ambulance	911
Gulf Coast Medical Center	850-769-8341
Admissions	850-747-7918
Financial Counselor	850-747-7914
Labor and Delivery	850-747-7700
Healthy Start 2101 Northside Dr. Unit #101, Panama City, FL 32405	850-872-4130
Medicaid Office 2505 W. 15th Street, Panama City, FL 32401 Toll free	850-872-4185 866-762-2237
Quest Diagnostic Laboratory	
Panama City	866-697-8378
Lynn Haven	850-360-1001
Panama City Beach	850-233-6420
Marianna	850-482-4918
Port St. Joe	850-532-6034
WIC/Nutrition 597 W. 11 th St. Panama City, FL	850-872-4666



Community Services

Service	Agency (ies)	Address	Telephone Number
Adoption	1. Catholic Charities of Panama City	1. 3128 11th St. Panama City, FL	1. 850-763-0475
	2. Children's Home Society 3. Adoption Center	2. 704 W. 23rd Street, Bldg E-40 Panama City, FL 32401	2. 850-747-5411
	4. Florida Baptist Children's Home	3. 3 Clifford Drive, Shalimar, FL4. 8415 Buck Lake Road, Tallahassee,	3. 850-785-0108 4. 850-878-1458
		FL 32317	
AIDS and HIV	1. BASIC 2. Bay County Health Department	1. 432 Magnolia Ave Panama City, FL 2. 597 W. 11 th St. Panama City, FL	1. 850-785-1088 2. 850-872-4455
Breastfeeding Support	1. Healthy Start Lactation Co. 2. Breastfeed Bay Hotline	nsultant	1. 850-215-1320 2. 850-747-5775
Child Medical Services/	1. Bay County Health Department	1. 597 W. 11th St. Panama City, FL	1. 850-872-4455
	2. Children's Medical Society	2. 914 Harrison Ave, Panama City, FL	2. 850-747-5411
Clothing/ Baby items	1. Catholic Charities of Panama City	1. 3128 11th St. Panama City, FL	1. 850-763-0475
	2. Children's Home Society	2. 704 W. 23rd Street, Bldg E-40 Panama City, FL 32401	2. 850-747-5411 3. 850-785-1721
	3. Family Service Agency 4. The Pregnancy Resource Center	3. 114 E. 9 th St., Panama City, FL 4. 701 Jenks Ave., Panama City, FL 32401	4. 850-763-1100
	5. A Sheltering Tree	5. 1616 Cincinnati Avenue, Panama City, FL 32401	5. 850-763-1433
Counseling	1. Life Management	1. 525 E 15 th St., Panama City, FL	1. 850-522-4485
Mental Health Services	2. Florida Therapy Services	2. 459 Grace Ave., Panama City, FL 32401	2. 850-769-6001
Drug Abuse	1. CARES	1. 4000 E. 3 rd St. Panama City, FL	1. 850-872-7676
Treatment	2. Life Management	2. 525 E 15th St. Panama City, FL	2. 850-522-4485

Community Services



Service	Agency (ies)	Address	Telephone Number
Early Child Development	1. Early Learning Coalition of NW Florida	703 15th St. Suite A, Panama City FL 32401	850-747-5400
Food Assistance	1. Catholic Charities of Panama City	1. 3128 11th St. Panama City, FL	1. 850-763-0475
	2. Beach Care Services	2. 9128 Front Beach Road, Panama City Beach, FL 32407	2. 850-235-3002
Food Stamps/ AFDC/WIC	1. WIC Bay County	1. 597 W. 11th St. Panama City, FL	1. 850-872-4666
AFDC/WIC	2. Bay County Dept of Children and Families	2. 2505 West 15 th St. Panama City, FL 32401	2. 850-872-7648
Housing	1. Panama City Rescue	1. 609 Allen Ave Panama City, FL	1. 850-769-0783
	Mission 2. Domestic Violence Program	2. Unlisted.	2. 850-763-0706
	3. Hidle House		3. 850-784-1020
Prescription Assistance	1. Bay County Health Department	1. 597 W. 11th St. Panama City, FL 32401	1. 850-215-0616
	2. Pancare	2. 2309 E. 15th St. Panama City, FL 32401	2. 850-329-4725
Teen	1 Duognon or Dogovino	1, 701 Jenks Ave., Panama City, FL	1. 850-763-1100
Services/	1. Pregnancy Resource Center	32401	1. 850-763-1100
Pregnancy Prevention	2. Children's Home Society	2. 704 W. 23rd Street, Bldg E-40 Panama City, FL 32401	2. 850-747-5411
	3. Hannah's House Maternity Home		3. 850-784-1020
	4. The Bridge Transitional Living Program		4. 850-481-8501
Transportation	1. Medicaid Transport		1. 850-785-0808
	2. Bay Town Trolley	2. Multiple locations	2. 850-769-0557



Providers

At Emerald Coast OB/GYN we provide specialized care to the pregnant woman and her family. Our providers practice as a team to give the very best of medical and nurse-midwifery care to the women we serve.

Henry Breland, George Ramie, Samuel Wolf, Toni Pennington, Kimberly Bain, Justo Maqueira, Jacob Martin and Heather Pate are our obstetrical physicians. Amanda Rodriguez, Shawn Brooking, Kim McConnell, Shelrel Berry and Cristina Stevenson are our certified nurse-midwives. All are board certified by their professional certifying organization and all are licensed in their respective field by the state of Florida. Our combined experience in the field of obstetrics exceeds 175 years! Together, we provide continuous coverage of all obstetric services to all of our patients at every hour of every day - throughout the year.

Here is some brief information about our providers. For a more complete bio, please refer to the staff section of our website.

If you desire a midwife to deliver your baby, our practice will make every attempt to accommodate your request. Simply notify our check out staff at the end of your appointment and they will have you sign our "Request for Certified Nurse Midwife Care" form and your chart will be noted.

Henry Breland, MD

Dr. Breland was born and raised in Marianna, Florida, He is the fourth of five children. His father, Jabe Breland, was a physician who practiced in Marianna for over forty years. His mother, formerly Betty Baker, was born and raised in Panama City. He attended Birmingham-Southern College for two and a half years and then transferred to Florida State University into the PIMS program, a forerunner to the medical school that is now there. In 1981 he received his Bachelor of Science degree in biology and transferred to the University of Florida to complete medical school, graduating in 1984. From there he did a year of internal medicine in New Orleans, Louisiana at Tulane University, and this was followed by a year of general surgery training in Huntington, West Virginia at Marshall University. For the next several years he worked in ambulatory care and emergency medicine in West Virginia until entering ob-gyn residency in 1993 in Charleston, West Virginia. After graduation in 1997 he remained in West Virginia for four years practicing ob-gyn. He then relocated to Panama City in 2001 to join the practice here at Emerald Coast OB-GYN. He is married to the former Nancy Holley, also of Marianna. Together they have five sons.

George B. Ramie, DO

Dr. Ramie is a 1996 graduate of Bay High School. He then graduated with a Bachelors in Science in Biology at Florida State University. He graduated with his Doctorate from Nova Southeastern University with Honors. While at Nova, Dr. Ramie was distinguished as an Anatomy Fellow, and was involved in Academic teaching. He went on to complete an Internship and Residency in Obstetrics and Gynecology in the Henry Ford Health System in Detroit, Michigan. While in Residency, he completed the requirements for the Consortium in graduate medical education offered through the Michigan State University. There he was a Chief Resident of his graduating class. Dr. Ramie is Board Certified in Obstetrics and Gynecology by the American Board Osteopathic Obstetrics and Gynecology.

Dr. Ramie has a special interest in the diagnosis and management of high risk conditions affecting pregnancy. In Gynecology, Dr. Ramie's scope of practice includes Well Women care through all stages of life. He has special interest in care for women in the perimenopausal and menopausal periods, particularly as it relates to aging. He embraces a holistic approach, utilizing Bio identical hormones to restore hormone levels. His scope of practice also include pelvic floor reconstructive surgery, vaginal surgery, advanced Laparoscopic surgery, daVinci Robotic surgery and the general management of most gynecologic conditions.

Dr. Ramie has three children and makes his home in Panama City, He is passionate about the delivery of a high quality health care experience in the office and at the hospital. Dr. Ramie is involved in medical staff leadership, serving on various panels and boards that assure safety and quality for patients. He strongly believes that care should be patient driven through a non-judgmental, empathetic, and caring relationship that puts the patient-physician relationship first.

Samuel Wolf, DO

Dr. Samuel Wolf is a native of Panama City. He received a B.S. in biology at Florida State University. He earned his doctorate (D.O.) at Nova Southeastern University College of Osteopathic Medicine. He completed a traditional rotating internship through the Michigan State University Health System, where he was awarded Intern of the year. He did his ob/gyn residency at the University of Florida Health Science Center / Shands Jacksonville. During his residency, Dr. Wolf served as administrative chief resident and was the recipient of many awards for teaching and research, including the prestigious Jelks Award for outstanding resident clinician of the year. He has had extensive experience in clinical gynecology and minimally invasive robotic gynecological surgery. He joined Emerald Coast OB/GYN in August of 2005 and received his board certification from the American Board of Obstetrics and

Gynecology in December of 2007 and is a Fellow of the American College of Obstetricians and Gynecologists. He is a published author of peer reviewed research in Obstetrics and Gynecology and on the board of directors for Florida Woman Care, a group of over 300 OBGYN's in the state of Florida. He has also served as the President and Medical Education Chairman for the Emerald Coast Medical Association along with various positions in leadership at both the state and national level. He served as Chief of Staff for Gulf Coast Medical Center 2016-2017. He is also the author and editor of 10 editions of this book. In his off times, Dr. Wolf enjoys spending time with his family and loves kitesurfing, snowboarding and hockey. He also started "The Vagicians" educational podcast in 2021 — listen in at www.wwthevagicians.com

Toni Pennington, MD

Dr Pennington joined Emerald Coast Obstetrics & Gynecology in 2008. She was born and raised in Sykesville, Maryland. She received her Bachelor of Science degree from Virginia Polytechnic Institute and State University and her medical degree from the Medical College of Virginia. She completed her internship and residency at the University of Florida Health Science Center/Shands-Jacksonville where she received multiple teaching awards, including the Berlex Teaching Award. She is board certified by the American Board of Obstetrics and Gynecology and is also a fellow of the American College of Obstetricians and Gynecologists.

Since arriving in Panama City, she has served in various board positions and currently serves as president of the Emerald Coast Medical Association. She is also the current OB GYN department chair at Gulf Coast Regional Medical Center. She particularly enjoys working with expectant mothers, as well as teens and their unique health care needs. She is also trained in minimally invasive robotic surgery.

In her spare time, Dr Pennington enjoys spending time with her husband, Scott, and their two boys. She also enjoys painting, reading, scuba diving, fishing and hiking.

Kimberly W. Bain, MD

Dr. Bain was born in Anniston, AL and grew up in Atlanta, GA. She earned a B.S. in Biology from Indiana University in 1986, graduating Summa Cum Laude and Phi Beta Kappa. She attended Vanderbilt University School of Medicine in Nashville, Tennessee, receiving her M.D. in 1990. She completed an OB/GYN residency at the University of Tennessee in Memphis. After residency she remained on the faculty for 2 years as an Instructor, teaching residents and medical students as well as seeing patients in an office setting. Dr. Bain then moved to Anniston, Alabama, where she was in private practice for 10 years. From 2006 to 2010, she was in private practice in Lexington, Kentucky where she also served as an Instructor for the University of Kentucky

Department of OB/GYN. In 2010, Dr. Bain and her family relocated to Panama City where she joined Emerald Coast OB/GYN. Dr. Bain has maintained Board Certification in Obstetrics and Gynecology since 1997 and is a fellow of the American Congress of Obstetricians and Gynecologists.

She and her husband, who is also a physician, met and married while they were at Vanderbilt Medical School. They have a son and a daughter.

Dr. Bain enjoys cooking, traveling, gardening, fishing, stand-up paddleboarding, SCUBA diving, and practicing her college French.

Dr. Bain has a special interest in fitness and nutrition. She has identified techniques and knowledge that will help women of all ages remain healthy and fit. She reached a point of frustration in her own fitness journey, which she successfully addressed using specific dietary and exercise routines. She understands the challenge of fitting a commitment to healthy lifestyle and fitness into a hectic schedule. But she is passionate about preventive medicine and the huge benefits that develop, and she truly enjoys working with her patients to help them feel good, look good, and be their healthy best!

Justo Maqueira, Jr., MD

Dr. Justo Maqueira, Jr. was the first physician trained and approved to perform daVinci Robotic Gynecologic procedures in the region. Since his first daVinci Robotic hysterectomy in February 2007 he has completed over 800 procedures using the daVinci Robotic Surgical System, making him the most experienced gynecologic daVinci surgeon in this region. Currently Dr. Maqueira is the only gynecologic surgeon in the region qualified to Proctor other surgeons with Intuitive Surgical on the daVinci Robotic Surgical System. Since proctoring his first case in July 2007 he has instructed over 100 Gynecologic Surgeons using the daVinci Robotic Surgical System around the country. He has a special interest in complex minimally invasive gynecologic surgery and is Board Certified by American Board of Obstetrics & Gynecology.

Jacob Martin, MD

Dr. Jacob Martin is a native of Panama City, born and raised in our small town. He received a B.S. in Medical Science at Southern Adventist University in Tennessee. He was then off to California where he earned his doctorate (M.D.) at Loma Linda University School of Medicine. He came back to the south to complete his OBGYN residency at the University of Kentucky. While there he was awarded the Award for Excellence in Female Pelvic Medicine and Reconstructive Surgery, The Award for Excellence in Minimally invasive Gynecologic surgery, and the Resident of the Year awarded by the Society for Academic Specialists in General Obstetrics and Gynecology. He has a special interest in gynecologic surgery and is passionate about giving his patients outstanding and individualized care.

Outside of medicine he enjoys spending time on our beautiful Gulf: scuba diving, fishing, kayaking, anything he can to get out on the water! While all of that is wonderful, his favorite thing is to spend every moment he can with his beautiful wife and new daughter. He feels so blessed to be raising his family in this beautiful place we get to call home.

Heather Pate, MD

An Alabama native, Dr. Pate graduated from Alabama College of Osteopathic Medicine in 2018 and completed her residency at Oklahoma State University in 2022. During her education, she also traveled to rural Jamaica with the CAMEO organization to provide medical supplies and assist physicians in treating patients who were unable to obtain regular healthcare secondary to geographic and financial disadvantages. Dr. Pate, her husband Blake, and their 2-year-old daughter are happy to call Panama City home now. She is excited to be working in the office where she completed her clinicals and is eager to give back to the community she learned so much from. In her free time, Dr. Pate enjoys spending time outdoors with her family and sweet husky, Laela. She also enjoys working out, hiking, traveling, and Alabama football!

Michelle Lang MSN, APRN-C

Michelle grew up in Jacksonville, Florida and moved to Panama City in 1995. She is a graduate of Gulf Coast Community College and Florida State University. She also received a Dual Master's Degree from the University of South Alabama as a Women's Health Nurse Practitioner and Women's Health Clinical nurse. She takes special interest in adolescent gynecology as well as traditional and bio-identical hormone replacement therapy. She has worked in the specialty of women's health since 1999. Michelle joined Emerald Coast OB/GYN in 2003. She holds National Certification as a Women's Health Nurse Practitioner.

Melisa Summers MSN, APRN-C

Melisa was born and raised in Panama City and joined Emerald Coast OBGYN in 2006 as an LPN. While working for Emerald Coast, she graduated from Gulf Coast State College with an Associates of Science in Nursing and also earned her Bachelor's degree in nursing from Chamberlain College of Nursing. In 2015, she left Emerald Coast to begin her career as an RN and began working at Bay Medical Sacred Heart. She returned to school to finish her Masters of Science in Nursing from Frontier Nursing University, and returned to Emerald Coast OBGYN in 2020 to follow her passion of providing exceptional care and education to women during all stages of life. She is board certified as a women's health nurse practitioner and is currently conducting our pelvic floor rehabilitation therapy program. Melisa is married with 3 children and enjoys spending time with her family, boating and spending time at the lake, and cheering on her kids at their sporting events!

Amanda Rodriguez MSN, APRN-C, CNM

Amanda spent the majority of her younger years along the Gulf Coast of Mississippi and Florida and is thrilled to have relocated to Panama City since August 2018. She is married and together they have a blended family of 10 and 1 grandson withe hope of many more grandchildren. She earned her Nursing Diploma at St. Elizabeth's School of Nursing in Lafayette, Indiana in 1998. She spent her early career in Pensacola, Florida, Oklahoma, and Georgia. From 2003 to 2018, she lived in Indiana and was a nurse in Labor and Delivery. During this time, she returned to school and graduated with her Master's degree in Nursing in 2010 from Indiana University. She practiced as a Nurse Practitioner from 2010 to 2014 at Healthnet in Indianapolis, IN. She is certified through NCC as a Women's Health Nurse Practitioner. She quickly realized that she missed the miracle of birth and could not wait to help women through this process again. She returned to school and received her Post Master's in Midwifery from University of Indianapolis in Indiana in 2013 and has practiced as a certified nurse-midwife since 2014.

Shawn Brooking DNP, APRN, CNM

Shawn's passion for women's health began as a child influenced by her grandmother who was a nurse in Brooklyn. After completing her bachelor's degree from California San Bernardino, she followed her passion and became a labor and delivery nurse for the U.S. Navy. After leaving the Navy, Shawn sharpened her skills as an L & D nurse in Southern California where she felt the desire to provide more care to her patients as a midwife. Shawn completed her Master's in midwifery from the Frontier School of Midwifery in 2010 and continued her work in Southern California. In 2014, her drive as a provider in women's health and midwifery pushed her to earn her Doctorate in Nursing Practice, also from Frontier University. Since completing her DNP, Shawn has worked in both North Dakota and previously in Florida. Now "back home" in Panama City, Shawn enjoys running and spending time near the water with her family. Shawn looks forward to being your provider in women's health, labor, and delivery.

Kim McConnell MSN, APRN, CNM

Kim earned her Masters of Science in nursing with focus on midwifery from Frontier Nursing University in 2018. She worked as a medical/surgery nurse for 2 years, a labor and delivery nurse for 3 years, and a nurse midwife for almost 2 years. She is originally from Leesburg, GA but has lived in the North Georgia Mountains and Florida previously. Kim and her family moved to Panama City for a change in lifestyle and are excited to call it their new forever home! She has future plans to return to school for her Doctorate of Nursing Practice and Women's Health Nurse Practitioner certificate. Kim has 3 grown children, lots of pets, and has been married for 24 years. She enjoys going to the beach (one of the main reasons that brought her back to Florida), saltwater fishing, boating, kayaking, and camping. Kim is passionate regarding her desire to provide the very best evidence-based care to all women with shared decision making for their care throughout their lifetime.

Shelrel Berry MSN, APRN, CNM

Shelrel was born in a small city in rural Alabama, the daughter of a farmer and a homemaker, and seventh of nine children. Shelrel acknowledges that her love for midwifery started at an early age when she was introduced to a lay midwife when she was just an adolescent. This sparked an interest in nursing, particularly, labor and delivery nursing. At that moment she realized that it was God's plan for her to pursue nursing and subsequently midwifery as a career and what she refers to as her ministry. She believes that midwifery is a gift given to her to be "with women" encouraging and empowering them through one of the most miraculous moments in life.

Shelrel received her Bachelor's of Science in Nursing in 1986 and Master's of Science in Nursing in 1994 from Troy University. She received a second Master's of Science in Nursing as an Advanced Nursing Practitioner with a Specialty in Midwifery from the University of Alabama at Birmingham in 1996. She has worked as a registered nurse primarily in Maternal Child Nursing in the Labor and Delivery, Post-partum, and Nursery Units. After completing her Midwifery Certification in 1997 she began her midwifery career in Troy, Alabama and has been providing midwifery care for over 22 years throughout south and southwest Alabama, and southwest Georgia.

Shelrel is married to Willie Frank Berry and they have four children and currently six grandchildren. Shelrel's husband is a veteran of the U.S. Army and retired police investigator. They are a family-oriented couple that enjoys serving God by serving others. Shelrel also enjoys traveling and spending time with family and friends. She is extremely excited and thankful that she is able to continue to provide women with caring, compassionate and comprehensive quality health care at Emerald Coast OBGYN.

Cristina Stevenson MSN, APRN, CNM

Cristina Stevenson is a Bay County native and board certified nurse midwife (CNM) who is passionate and committed to the excellent care of all women, whatever their needs might be—from preconception to delivery and beyond. She began her career as a volunteer at Gulf Coast Medical Center while dual-enrolled in high school and college. She earned her ADN from Gulf Coast State College in 2001 and went to work immediately in labor and delivery- a dream she had since childhood. She got her BSN from Jacksonville University and traveled the nation answering the call of her heart serving the women, babies, and families in her care.

In February 2023 she earned her Masters Degree in Midwifery from Frontier Nursing University in Versailles, Kentucky, completing her clinical training in Tallahassee, Florida, and Dothan, Alabama. She brings more than 20 years of experience working to facilitate the miracle of birth and the care of women and their babies.

When not at work, Cristina enjoys traveling, spending time with her husband and their three children, and their big busy family on the family farm.

As a Certified Nurse Midwife, Cristina brings a wealth of experience, a passion for life, and a commitment to excellence in the care of the whole woman.

Gulf Coast Medical Center



Emerald Coast OB/GYN chooses to use only Gulf Coast Medical Center for childbirth care. We think you will agree with us that GCMC is a very nice facility in which to give birth. * If you go to Bay Medical Center ER, you will NOT be able to be seen by your providers since they do not practice at that hospital. This obviously complicates continuity of your care so we encourage you to use the ER at Gulf Coast.

GULF COAST MEDICAL CENTER:

449 West 23rd Street. Panama City, FL 32405

Located off of 23^{rd} Street near the intersection of State Avenue. The Women's Center is located on the 2^{nd} Floor – closest to the main entrance.

Website: www.gcmc-pc.com/services/womens-center

Labor & Delivery:	747-7700
Accreditation:	Full
Average births per month:	180-200
Post Partum Rooms:	Private
Bathrooms:	Complete with soaking tubs.
Infant Rooming-In:	Yes
Well Baby Nursery:	
Lactation Consultant:	Full-Time
Childbirth Classes:	Yes, call 747-7700
Anesthesia Services:	In-House 24 hours.
Pediatricians:	Most local doctors are on-staff
Dedicated Maternal Fetal Medicine Doctor	Dr. Ramada Smith
Dedicated pediatric surgeons	Dr. Taylor and Dr. Walford
Obstetrical Operating Rooms:	2
Pediatric Surgical Subspecialties:	Yes

EMERGENCY CARE

Provided in the Emergency Room until the $20^{\rm th}$ week. After 20 weeks, stable patients will initially be evaluated in Labor & Delivery.

NEONATAL INTENSIVE CARE UNIT

A Level III neonatal unit is maintained at Gulf Coast Hospital and is available for babies requiring advanced care. The hospital contracts for the full-time services of a neonatologist. Occasionally, a baby is born with an unforeseen problem requiring very specialized care. At such times the situation is handled on an individual basis and the baby is transferred to the hospital where that particular special care is available.

PRE-REGISTRATION

You must pre-register with Gulf Coast Medical Center around the 28th week of your pregnancy. Please go to the following website to do this: https://gcmc-pc.com/

patients/registration.dot or you can just Google "Gulf Coast Hospital Preregistration". If you are paying for childbirth expenses "out-of-pocket", you will want to visit the Admission Office at the hospital in order to learn of special discounts they may offer for advanced payment. You can also ask our check out staff for a paper version of this to complete.

LACTATION SUPPORT

Lactation consultants are on staff to help you initiate breast feeding during your postpartum stay. Also, Gulf Coast Medical Center has a virtual, interactive breastfeeding webinar that is free of cost. Please visit their website (gcmc-pc.com) for dates and times.

HELPFUL BREASTFEEDING NUMBERS

Bay County Healthy Start — 850-215-1320 Breastfeed Bay Hotline — 850-747-5775 Franklin County Healthy Start — 850-370-6351 Gulf County Healthy Start — 850-227-1276 WIC Breastfeeding Hotline (24/7) — 850-747-5775 La Leche League — 850-525-3243

LOCK DOWN

If you need to go to Labor & Delivery between the hours of 9PM and 6AM you will find that the front entrance is locked for security purposes. In this situation, you will need to enter the hospital at the Emergency Room entrance located at the east side of the facility. Also, be sure to tell family and friends that incoming telephone service to patient care rooms is cut-off at 9PM to allow you to rest. You may still call out. The doors to the Women's Center are kept locked. All visitors need to be buzzed in and out of the unit by the receptionist. Also, visitors need to sign in at the front desk and wear a visitor sticker while in the Women's Center. The post partum unit is now located on the 3rd floor of the new tower at Gulf Coast Hospital. ***The mother/baby unit is NOT accessible to visitors via the labor unit. Visitors must go down stairs and around the hall to the ICU elevators.***

VISITORS

During labor, you will be expected to limit your visitors to no more than two at a time. During the time of delivery, you will be limited to keeping up to two people with you (no trading out) until you and baby are safe and sound during the recovery period. This is a hospital policy and exceptions can only be made by making special arrangements in advance with the nursing administrator responsible for the Labor & Delivery unit. The waiting room in the Labor & Delivery unit is small and usually crowded. Remember that there is a much larger waiting room located off the first floor lobby, next to the gift/snack shop. Visitors are also welcome to purchase food and beverages in the hospital cafeteria on the first floor. Once you have given birth and completed the immediate recovery period you may have visitors with more lenient limitations. Visitors will be expected to leave the building at 9 PM. One person may stay with you overnight. That person must be old enough to offer assistance to you. Family and friends that are ill should not come to the hospital. Please call the hospital for the most up to date visitor policy.

*If you would like a tour of Gulf Coast Medical Center's Women's Center, please call 850-747-7700.

Laboratory Tests



Towards the beginning of your pregnancy care, you need to review and understand the available lab tests. As technology offers more tests, counseling for these has become more complicated. This section of the booklet will attempt to break down the essential information in an easily understandable way. Testing guidelines and available tests change over time. This is not a comprehensive review. It is important to discuss any questions you have with your provider.

Typically, patients will have blood work performed early in the pregnancy and then again at 26-28 weeks for some routine follow-up tests. We are proud to be the only obstetric practice in the area to offer this service right in our office at no extra cost to you so that you do not need to have blood work performed at an outside collection site, such as Quest or LabCorp. Please be patient with our blood collection staff. Our phlebotomists are highly skilled and can get very busy. The lab specimens are collected by us and transferred to the lab preferred by your insurance or to the lab of your choice if you are uninsured. Most insurance companies in our region prefer to use QUEST Labs. However, if your insurance requires that your specimens be sent to another lab for processing, please let our phlebotomist know before your blood is drawn. For patients without insurance, please inquire about our reduced fees negotiated with Florida Medical Lab.

There are certain labs that will need to be done through Gulf Coast Medical Center. This includes the GBS culture done at 35-37 weeks. This specimen will be collected in our office and sent to Gulf Coast Medical Center's laboratory for you (see page 60 for more information on GBS).

Standard Tests

Routine initial tests include the following: a complete blood count, blood type, Rh, antibody screen, Rubella immunity screen, RPR (syphilis screen), Hepatitis B and C screening, HIV test, thyroid screen, and a urinalysis with culture. An early diabetes screen will be ordered if you are overweight based on your height. A Pap smear will be performed if indicated at your first exam. Pap smears are generally not indicated for patients less than 21 years of age. Gonorrhea and Chlamydia testing will be performed on all pregnant patients unless you specifically decline. If you have never had chicken pox or the vaccine, testing for immunity will be necessary with your prenatal labs. We do not routinely test for genital herpes, so please let us know if you have ever been exposed or diagnosed with it.

Around 27-29 weeks, a repeat complete blood count, RPR (syphilis screen) and a 1-hour gestational diabetes screen will be ordered. Once again, this can be done in conjunction with your OB visit. You DO NOT need to be fasting for the gestational diabetes test. However, eating foods high in carbohydrates just before the test will increase your risk of having an elevated diabetes screen (or a false positive test). Examples of foods that usually will not alter the results are eggs, meats, and cheeses. Water is always OK. You should plan to spend about 1 ½ hours at our office or at the lab to complete this test.

3 Hour Glucose test

This test will only be ordered if your 1-hour gestational diabetes screen is elevated. If this test is ordered, you will need to schedule it in advance at the lab of your choice (Quest or LabCorp). You should plan to budget about 4 ½ hours and you will be required to stay within the testing facility for the duration of the test. Bring a good book, and iPad, or something else to do while waiting. Eat a balanced diet that contains at least 150 grams of carbohydrates per day for 3 days before the test. Fruits, breads, cereals, grains, rice, crackers, and starch vegetables such as potatoes, beans, and corn are good sources of carbohydrates.

This is a fasting test. Do not eat, drink (except water), smoke, or exercise strenuously for at least 8 hours before your first blood sample is taken.

Tell your provider about all prescription and non-prescription medications you are taking. You may be instructed to stop taking certain medications before the test.

(Instructions from the American College of Obstetricians and Gynecologists and the American Diabetes Association)

Optional Testing

The American College of Obstetricians and Gynecologists recommends that we offer all patients screening for Cystic Fibrosis, Spinal Muscular Atrophy, and Hemoglobin disorders such as Sickle Cell Disease. There are other genetic conditions that are more prevalent in certain ethnic groups, particularly people with Ashkenazi Jewish ancestry, where expanded genetic carrier screening may be indicated (Additional patient information at acog.org/Carrier Screening). Traditional testing for fetal problems includes:

Cystic Fibrosis (CF) Carrier Screening

Cystic Fibrosis is one of the most common genetic disorders in the United States. It is associated with breathing problems, recurrent lung infections, as well as problems with digestion. There are medications that can be given to help but even with treatment, this condition causes life-long illness and a shortened life expectancy. Cystic fibrosis does not affect intelligence or physical appearance.

Cystic fibrosis is a genetic disorder that results from mistakes or changes to a certain gene. All genes come in pairs, one from the mother and the other from the father. Cystic Fibrosis is present only when an individual inherits two changed (abnormal) genes, one from each parent. When an individual has one changed gene and one normal gene, the person is a carrier. No significant health problems are caused from being a carrier of cystic fibrosis. Unless someone had testing, they will never know they are a carrier. The chance an individual is a carrier depends on their family history and ethnic background. You can be a carrier even if no one in your family has Cystic Fibrosis. Once you are tested for this, you will never need this test again. All babies are tested for CF at birth.

Spinal Muscular Atrophy (SMA)

Spinal Muscular Atrophy is a disorder that results from a missing or defective SMN1 gene. When this gene is missing or not working properly, the body cannot make enough of a particular protein that is needed for motor neuron cell survival. Everyone is born with a certain amount of these motor neuron cells. These cells are responsible for communicating with the muscles to tell them to work properly. When the body cannot make enough of this protein, the motor neuron cells become weaker and weaker over time and will eventually stop working altogether. They will lose all function and the cells will die. As a result, muscle function deteriorates and normal functions such as breathing, eating, speaking, and lifting the head, become difficult. SMA does not affect cognition, emotional development, learning, academic, or sensory ability.

Hemoglobin Disease Testing

This optional test will assess genetic problems with the mother's hemoglobin, such as Sickle Cell Disease or Thalassemia. We strongly recommend that African American patients and patients with Mediterranean ancestry be tested for hemoglobin disease unless they have previously tested negative.

Cell-free Fetal DNA (cfDNA) testing

Now that the baby's DNA can be tested by a maternal blood test, amniocentesis is rarely needed. Most insurance companies including Medicaid will allow patients over 35 to bypass traditional testing and to go straight to the fetal DNA blood test. This test is the most accurate non-invasive screening test for Down Syndrome, but false positive and false negative results can still occur. Younger patients are at a higher risk of having a false positive test. The test can usually also determine the sex of the fetus and can be performed after 10 weeks. Amniocentesis is still recommended to verify a positive result. This test does not detect all chromosomal disorders and sometimes the result can be inconclusive meaning neither positive nor negative. This

test is available to all our patients regardless of insurance coverage, but you may incur additional cost.

AFP

This is a simple test for neural tube defects such as spina bifida that is collected after 15 weeks unless you decline.

The QUAD Screen

This is a blood test that will screen for Down syndrome, and other genetic problems, plus neural tube defects such as spina bifida and is performed at 15 weeks. It is important to understand that the QUAD screen has a high number of "false positive" results. If your test is positive, IT DOES NOT MEAN, there is something wrong with your baby. It means additional testing needs to be offered to you.

Unfortunately, CF, SMA, and Hemoglobin disease are not the only genetic problems carried. There are many more that we do not specifically list on our lab consent such as Fragile X disease and certain types of Muscular Dystrophy. It is for this reason that giving your provider a detailed and accurate family history is extremely important. If you know of a disease or condition that runs in your family, or even if you have a family member with an unexplained form of mental retardation, please let us know so that we can offer you additional testing.

Ultrasound



With our basic ultrasound we can determine due dates, number of babies, placental location, amniotic fluid volume, and basic anatomy. While our scans are not designed to detect specific fetal anomalies, if we suspect problems during the course of an ultrasound, we will offer referral to a perinatologist (MFM) where a higher level ultrasound designed to detect specific fetal anomalies will be performed. All patients have the option of being referred for a higher level ultrasound. Notify your provider is you want this and we will schedule it for you.

SCREENING ULTRASOUND

We offer in-office ultrasound as a service to our patients. The anatomy screening ultrasound will typically be done around 20 weeks. These ultrasound images will be read by a board certified maternal fetal medicine high risk OB doctor. This ultrasound is useful in determining the size, position, and internal anatomy of the fetus, but there are limitations. Some abnormalities may not be detected until later in the pregnancy, or not at all. Ultrasound performed in the second trimester of pregnancy on obese mothers is less likely to detect structural anomalies or other markers of chromosomal disease in the developing fetus. Fetal sonography is not sensitive to detect all fetal abnormalities, structural of functional, and a "normal" exam should not be assumed to mean that the fetus is without anomalies.

OPTIONAL AND NON-COVERED SCANS

We offer 4-D ultrasound and fetal sex determination scans. While medically indicated ultrasounds are generally covered by insurance with specific indications, sex determination and 4-D ultrasound for bonding with the fetus are not covered by insurers. We offer two options if you are interested in having ultrasound beyond those which are covered by insurance plans.

GENDER DETERMINATION (NON 4-D) (\$75.00)

This is performed in 2-D mode and is a very limited survey of your baby with specific focus on the genitalia. You are more than welcome to film a short video from your smart phone. You will also be given photo images at no additional cost. Regarding sex determination, we strive to be accurate and have a very good track record, but cannot offer a 100% guarantee!

4-D ULTRASOUND (\$150.00) (\$75 redo)

Our ultrasound machines have the latest 4D technology to allow real time 3 dimensional images of your baby. You are more than welcome to film a short video from your smart phone. The best time for 4-D ultrasound is between 30 and 34 weeks gestation. ***Please remember that not all babies cooperate during their ultrasounds and we can offer no guarantee that images and photos will be "picture perfect". In the event your 4-D ultrasound pictures are not as good as you hope, we will gladly repeat your ultrasound at the discounted price of \$75.00. Both, the gender determination and 4-D ultrasound must be scheduled in advance and can be arranged at the Check Out window. Please note that your 4-D scan will take longer than any of your other ultrasounds and as a result, will not be scheduled on the same day as a visit with your provider. They will only be scheduled when you are between 30 and 34 weeks gestation.



Approved Medications

While most medications now carry "pregnancy warnings" the medications listed on these two pages are "over-the-counter" and are those we are very comfortable using during pregnancy. We believe the medications on this list will not cause harm to your unborn baby.

CONSTIPATION

• Surfak, Colace, Metamucil, Fibercon, magnesium oxide, or any overthe-counter stool softener may be helpful if used according to package directions. Avoid use of laxatives.

COUGH & CONGESTION

• Dimetapp, Mucinex, or Robitussin – use according to package directions.

DIARRHEA

• Imodium AD. Use according to package label.

GAS PAIN OR FLATULENCE

• Mylicon or simethicone products – use according to package directions. Also, avoid drinking through a straw. You may also use Beano before meals.

HEARTBURN, ACID INDIGESTION, OR GASTRIC REFLUX

• Gaviscon, Tums, Rolaids, Mylanta, Pepcid AC, Prilosec OTC etc. Use according to package directions.

HEMORRHOIDS

 Anusol products, Preparation H products, Tucks – use according to package directions.

APPROVED VACCINES

• The influenza (flu) shot and Tdap (tetanus, diphtheria, and pertussis) shot are safe to take during pregnancy and are highly recommended. See page (32) for details about these two vaccines. Be sure to get both of these shots to protect you and your baby!

INSOMNIA (SLEEP AIDS)

• You may use Tylenol PM, Unisom or plain Benadryl according to the package directions to occasionally help you get to sleep.

MILD DISCOMFORTS (HEADACHES, BACKACHES, "AROUND LIGAMENT" PAIN, ETC.)

• Tylenol – use according to package directions. We recommend that you avoid the use of ibuprofen, Motrin, Advil, or aspirin products **unless** specifically recommended by us. 400 mg of over-the-counter magnesium oxide works well to prevent migraine type headaches.

NAUSEA

- Emetrol. Use according to package label.
- Ginger capsules, 250mg. Take by mouth, every 6 to 8 hours on a regular basis. Peppermint can also help.
- Vitamin B-6 (25 mg tablet form). Take one tablet by mouth, every 6-8 hours on a regular basis.
- Unisom (doxylamine) 25mg <u>tablets</u>. Break them in half and take ½ tablet by mouth, every 6 to 8 hours on a regular basis. At this dose the remedy does not often increase sleepiness.
- Dramamine (dimenhydrinate). Take 50 to 100mg by mouth every 4 to 6 hours. Do not take more than 200mg a day if you are also taking Unisom.

SINUS CONGESTION (COLD OR ALLERGY SYMPTOMS)

- Chlortrimeton, Claritin, Zyrtec, or any Tylenol products, like Tylenol Cold, Tylenol Sinus, etc. Use according to package directions.
- You may use nasal saline spray as often as you desire.
- If you are running a fever (100.5) or have yellow/green drainage, call for an appointment.

SORE THROAT

• Throat sprays, lozenges – use according to package directions. Warm saline gargles are very helpful and can be used as often as you desire. If you have a fever of 100.5 or more, or exposure to someone with "strep throat", you need to call our office."

SUNBURN PROTECTION

• Be very careful about sun exposure during your pregnancy. Pregnancy hormones make it much more likely that you will burn very quickly. Use clothing, hats, umbrellas and UVB/UVA sun block to protect yourself. Also, remember to drink lots more water while outdoors in the heat. You will become dehydrated much faster while pregnant. This will cause uterine cramps.

VAGINAL ITCHING

• Monistat, Mycelex, or Gyne-Lotrimin. Use according to package directions.

*** Questions about any of the above medications or about any medications that are not listed should be discussed at an appointment <u>or</u> by by sending us a message on the portal. You may also call our office to speak with a member of our staff at 850-769-0338.



Prenatal Education

Perhaps at no other time in your life will education about your condition become more important to you and to those you love. Pregnancy is a brief but physically and emotionally challenging time in any woman's life. It's a great time to get yourself and your family prepared for the immediate future. Fortunately, many options and offerings are available to you!

For beginners, we recommend the book, <u>Your Pregnancy Week by Week</u> by Dr. Glade Curtis. You will find this book in most bookstores or at the public library. It can easily be purchased on-line at Amazon.com. Other helpful resources are listed at the end of this book.

Please note if you watch labor or birth shows on television, remember that the producers of these shows are generally filming the most sensational stories they can find and usually don't have patient education in mind during their production. SCARY!

CLASSES

Community Pregnancy Information Classes: The Pregnancy Resource Center offers a variety of prenatal and child rearing classes. The classes are provided at no charge and have the added benefit of special coupons that can be redeemed for a variety of essential supplies and equipment that expectant mothers might need. To find out more about these classes please call: 850-763-1100.

Gulf Coast Medical Classes: Call the Women's Center at 850-747-7700 to inquire about any additional childbirth education classes that may be available at the hospital.

Morning Sickness



Nausea and vomiting with pregnancy, often referred to as "morning sickness" affects 70 to 85% of all women. It is caused by an increase in hormones released by the developing placenta early in the pregnancy. Many women experience only a queasy stomach or loss of appetite. Some women find it to be a debilitating condition. "Morning sickness" may occur at any time during the day or night, with symptoms usually starting between the 4th to 7th weeks of pregnancy and ending by the 16th week for the vast majority (90%) of women. On rare occasions, nausea and vomiting may persist throughout the pregnancy.

WHAT CAN I DO?

No one remedy will be effective at all times. It usually takes a combination of many strategies to keep problems to a minimum. Here are some helpful tips.



- Use a vitamin that contains only folic acid, B6 & B12. Take it in the late afternoon or before bed with a small meal.
- Eat small amounts of appealing food every 2 hours. Be sure to eat proteins such as nuts, eggs, yogurt, cheese, meat, & fish.
- Use aroma & taste therapy. Use lemon drops, peppermint, pickled sushi ginger, etc.
- Sip iced beverages such as lemonade, water, or ginger ale.
- Nibble crackers, toast, or gingersnaps just before getting out of bed.
- Wear seasick prevention wrist bands around-the-clock. They cost less than \$10.00 and can be found in many drug or boat supply stores.

HERBS & MEDICINES

Studies have shown the following medications and herbal remedies to be helpful. If taken on a regular basis they may <u>prevent</u> nausea & vomiting – rather than to immediately relieve a current episode. All are available without a prescription.

- Ginger capsules, 250mg. Take by mouth, every 6 to 8 hours on a regular basis.
- Vitamin B-50 Complex (<u>tablet</u> form). Break them in half and take ½ tablet by mouth, every 6 to 8 hours on a regular basis.
- Unisom (doxylamine) 25mg <u>tablets</u>. Break them in half and take ½ tablet by mouth, every 6 to 8 hours on a regular basis. At this dose the remedy does not often increase sleepiness.
- Dramamine (dimenhydrinate). Take 50 to 100mg by mouth every 4 to 6 hours. Do not take more than 200mg a day if you are also taking Unisom.

PRESCRIPTION MEDICATIONS

If these suggestions do not help you - please contact us using the patient portal or give us a call. We can prescribe other medicines that are not available over-the-counter. Some of these medications are very expensive and some cause significant sleepiness which can disrupt your normal activities.

DIET AND DEHYDRATION

If you are having difficulty tolerating a regular diet, try a BRATT diet (bananas, rice, applesauce, toast, and tea). This diet is easy on your stomach and will help provide essential nutrients. If you cannot tolerate any solid food, Gatorade is helpful by providing calories and potassium.

If you are vomiting frequently and cannot keep clear liquids down for a prolonged period of time, please give us a call. On rare occasions patients require intravenous re-hydration in the emergency room or in the infusion center at Gulf Coast Medical Center.

Warning Signs During Pregnancy



The following list of symptoms can be associated with a serious problem with your pregnancy. If you experience any of the following, notify us by telephone immediately. We are on-call 24/7, 365 days per year to cover pregnancy related emergencies. Notifying us early, can allow us to prepare the hospital staff accordingly and help to avoid problems with delay in care.


- After 24 weeks, decreased fetal movement during any 12-hour period, especially when you are resting or during the evening hours.
- Fever of 100.5 degrees or higher.
- Vaginal bleeding.
- Persistent dizziness or fainting spells that occur even though you are well hydrated and well nourished.
- Persistent headaches that occur even though you are well hydrated, well nourished, and continue after taking Tylenol and resting.
- Sudden swelling of the hands and/or face.
- Blurred, 'sparkly', flashy, or strange vision. Little white floaters that occur occasionally and after rubbing your eyes are nothing to worry about.
- Pain or burning associated with urination.
- Sharp abdominal pain that persists after resting.
- Persistent vomiting and/or diarrhea despite using our recommendations in this booklets section on Morning Sickness (pg. 26) and Approved Medications. (pg. 23)
- More than 4 painful contractions per hour that do not respond to hydration and rest if you are less than 36 weeks. (see Pre-term Labor pg. 51)
- Rupture of the bag of waters or leaking of amniotic fluid. Please review our booklet section on ruptured membranes. (pg. 64)
- Persistent rash, such as redness or irritation of the skin.
- Pain under your incision if you have previously delivered by c-section.
- Any other symptoms that have recent onset and are concerning to you.





Nutrition During Pregnancy

Eating right during your pregnancy is one of the very best things you can do for yourself and your baby. Pregnant women are sometimes concerned about gaining weight during pregnancy. How much weight you plan to gain during pregnancy depends on your weight before becoming pregnant. Use our weight gain graph to keep up with the amount you gain as you progress through your pregnancy. Plotting your weight on this graph will help you keep your weight gain in perspective. Only weigh yourself 1 time per day in the morning after you void.



The Six Food Groups

Where Does Your Weight Go???

•	Blood	3.0 pounds
•	Breasts	2.0 pounds
•	Uterus	2.0 pounds
•	Baby	7.5 pounds
•	Placenta	1.5 pounds
•	Amniotic fluid	2.0 pounds
•	Retained tissue fluid	4.0 pounds
•	Stored fat/protein	7.0 pounds

Total: 29 pounds

EMERALD COAST DIETARY RECOMMENDATIONS

We recommend eating 3 meals and 3 small snacks throughout the day. The <u>South Beach Diet</u> can be used as an excellent guide for your diet. When using the <u>South Beach Diet</u> be sure to select foods from both Phase I and Phase II food groups. Avoid eating more than one starch or simple carbohydrate food at any one meal or snack. The majority of your carbs should come from fruits and vegetables. Remember that juices are mainly sugar, so try to dilute them with water or diet ginger ale for instance to decrease your carb intake.

Avoid all empty calorie and fast foods - as much as humanly possible! Other tips: Don't eat chips out of the bag, don't watch TV while you eat, try to have a small appetizer 30 minutes before meals, and try to make dinner the smallest rather than the largest meal of the day. **Sweet-tea and soft drinks are absolutely horrible for you during pregnancy. Please avoid them completely.**

Remember the most important foods for your developing baby are fats and protein.

Ideas for snacks: Nuts, avocado, cheese on whole grain crackers, yogurt, trail mix, protein bars, sunflower seeds, dried fruits, boiled eggs, raisins, pickles, and Edamame (soy beans) which can be found in the organic frozen food section.

EMERALD COAST RECOMMENDED FOOD SERVINGS

PREGNANCY FOOD PRECAUTIONS


- Moderate fried foods, and sugar intake. It's fine to use artificial sweeteners.
- Limit your use of caffeinated beverages to 2 servings a day.
- If you have lactose sensitivity it is fine to use lactose free products.
- Do not eat unpasteurized soft or hard cheeses. Check the labels.
- Seafood known to have high mercury content includes Shark, Swordfish, King Mackerel, Tuna and Tilefish. Canned light Tuna has less than Albacore. Limit these fish to one serving a week.
- Reheat prepared meats, such as hot dogs and deli meats reheat until steaming.
- Do not eat raw meats or seafood. Cooked oysters and sushi are fine.
- Excessive alcohol intake during pregnancy has been associated with numerous birth defects.

A healthy target weight gain for the average woman (BMI between 20 and 26) is between 25 and 35 pounds. If you were overweight at the start of your pregnancy (BMI above 26) your target should be to gain between 15 and 25

pounds. For thin women (BMI below 20) it is recommended that you gain between 28 and 40 pounds. Use the weight gain graph below above to keep up with the amount you gain (or lose) as you progress through your pregnancy.

Start by writing down your starting pregnancy weight on the bold line at the far left of the chart. Record your recommended target weight gain goal range along the bold line at the far right of the chart. As you continue through your pregnancy record any gains (or losses) that have occurred since you last weighed. Try to weigh on the same scale whenever possible and try to weigh with your shoes off, whenever possible.

Plotting your weight on this chart will help you keep your weight gain in perspective.



My Prenatal Progress

Date	Weeks	Weight	B.P.	Fetal Heart Rate	Uterine Fundal Height
		20			



Smoking

Smoking is very common in our community. In fact, 13.7% of all Americans smoke cigarettes and 3.2% of adult Americans vape. Nicotine can be as addictive as crank, heroin, or cocaine. It is always hard to quit, but the good news is that it's often easier to quit smoking during pregnancy. During pregnancy cigarettes often don't taste as good and pregnant women are more motivated to quit for their baby's sake! Cigarettes are particularly harmful to the developing fetus. Mothers who smoke are more likely than average to have the following problems:

- Miscarriage
- Small birth weight babies
- Stillborn babies
- Abruption (when the placenta separates from your uterus)

These problems arise because cigarette smoke causes premature aging of the placenta which is your baby's lifeline for oxygen and nutrition.

Problems with smoking are not limited to the pregnancy. Babies born to tobacco addicted mothers can have irritable symptoms from nicotine withdrawal. Babies living in homes where there is cigarette smoke are more prone to chronic lung problems such as asthma. Furthermore, these babies are at greater risk for sudden-infant-death syndrome. Finally, children are less likely to start smoking if their parents do not smoke. For all of these reasons (as well as for your own good health) it is extremely important to quit smoking. We will work with you and help you succeed! Our Health Care Navigator can assist you with the SCRIPT program that is designed to help you quit smoking within 7 days — ask your provider for more information.

TIPS FOR QUITTING GET READY

Set a quit date. Reduce the number of cigarettes you smoke. Get rid of ALL cigarettes and ashtrays in your home, car, and workplace. Don't let anyone smoke in your environment. Review your past attempts to quit (what worked and what didn't). Once you quit, do not smoke – not even one puff.

GET SUPPORT & ENCOURAGEMENT

Tell your family, friends, and coworkers that you are quitting and that you want their support. Ask them to not smoke in front of you or leave cigarettes out. Get individual, group, or telephone counseling. Avoid people who secretly want you to fail!!! Talk to us – we can help!!!

LEARN NEW SKILLS & BEHAVIORS

Distract yourself from urges to smoke. Talk to someone, take a walk, or get busy with a task. When you first quit, change your routines. Take a different route to work. Drink decaffeinated tea instead of coffee. Eat lunch outdoors. Do things to reduce your stress. Take a hot bath, exercise, read a good book. Take deep and relaxing breaths of clean air. Drink lots of water. Plan something enjoyable to do every day! Buy a yoyo, paddle ball, or something to occupy your hands.

GET MEDICATION & USE IT CORRECTLY

Medications can help you stop smoking and lessen the urge to smoke. Using one of these medications will practically double your chance of quitting! Ask us for advice about the use of these medications during pregnancy. While we do not prescribe nicotine patches during pregnancy, there are other effective medications that we can prescribe.

PREPARE FOR RELAPSE OR DIFFICULT SITUATIONS

Most relapses occur within the first 3 months after quitting. Use a relapse as a learning tool and move forward. Be aware of triggers like alcohol and being around other smokers who will ofter smoke with them.

ADDITIONAL SMOKING CESSATION RESOURCES:

Call Healthy Start @ 850-872-4130

Quitline @ 877-822-6669

"Vaping" is not safe during pregnancy. The contents of "vapor smoking" or "E-cigarette" products are unknown. Many of the products are imported from other countries and not tightly regulated by the FDA for safety.

The American College of Obstetricians and Gynecologists (ACOG) and the CDC both recommend against the use of e-cigarette products by pregnant and postpartum individuals, children, and adolescents, and adults who currently do not use tobacco products.



Important Immunizations During Pregnancy

INFLUENZA (FLU) SHOTS

Pregnant women and their babies are at increased risk of catching influenza (the flu). This is because your immune system is slightly diminished

during pregnancy, which increases your risk of infections. While influenza generally causes mild illness, pregnant women are more likely to suffer more severe symptoms and the flu can even trigger premature labor. Also at higher risk are newborns and any child under age 5. You need to take this very seriously. Flu (combined season & H1N1) shots are especially recommended during pregnancy. You and your other household members should be vaccinated just as soon as



vaccine becomes available. <u>Anyone</u> who will be in close contact with your baby should be vaccinated before your baby is born. *This is especially important because your new baby cannot receive influenza vaccine protection until she or he reaches 6 months of age.* There is evidence suggesting that vaccinated mothers can pass protection to the fetus and later to newborns through breast-feeding.

TDAP (tetanus, diphtheria, and pertussis) BOOSTER

America is experiencing a serious increase of Pertussis (whooping cough) illness in infants. Most infections are occurring in babies too young to have their own protection developed as they complete their DPT baby shots. Whooping cough illness can be life threatening to infants. Most deaths have occurred in infants younger than 2 months of age. The American Advisory Committee on Immunization Practices recommends that every pregnant woman be immunized with the Tdap vaccine between the 27th and 36th weeks of her pregnancy. Giving the vaccine late in pregnancy allows passive antibody protection to pass from the mother to her fetus. Optimal antibody protection begins occurring 2 weeks after the woman receives her immunization. Our office is able to administer the Tdap vaccine to our patients with commercial insurance. TriCare patients will need to receive their vaccinations at the base clinic. Our patients with Medicaid and those without insurance can receive their vaccine at the health department or walk-in clinics that carry the vaccine. Other household members and any other people expected to have close contact with the baby should also receive this vaccine before the baby is born.

Influenza (flu) Infection & Pregnancy



SYMPTOMS

Fever (usually high) with chills, headache, extreme fatigue, dry cough, runny or stuffy nose, muscle aches, sore throat, vomiting, and occasionally, diarrhea.

WHAT IF I GET SICK?

- Stay at home and limit contact with others.
- Get lots of rest.
- Treat any fever right away (use Tylenol.)
- Aggressively increase your fluid intake.
- Call our office as soon as you believe you have the flu symptoms listed above. We will review your situation and advise you if we think testing or anti-viral medication (i.e. Tamiflu) are needed. These medications work best if started within 48 hours of becoming ill. There is little information about the effect of these medications when used during pregnancy, but no serious side effects have been reported.




Influenza is spread from person-to-person by exposure to airborne viral particles and by touching surfaces that have been contaminated with secretions. It is not spread by eating pork! Here are some everyday preventive actions that can offer you protection:

- Avoid close contact with sick people.
- Cover your nose and mouth with a tissue, or your sleeve, whenever you cough or sneeze. Throw your used tissues into the trash and wash your hands.
- Wash your hands often with soap and water, especially after you cough or sneeze. Try
 to wash for 20 seconds (sing Happy Birthday for good timing!) If soap and water is
 unavailable, alcohol-based hand cleaners are also effective. They should be rubbed in
 until dry.
- Avoid touching your eyes, nose or mouth. Germs enter your body through these pathways.
- Make a plan for how to care for sick family members.
- Stock up on household, health, and emergency supplies, such as Tylenol, tissues, soap
 and alcohol-based hand cleaners, canned goods such as soups and other non-perishable
 foods, etc.
- Call us if you have been exposed to someone with a flu-like illness. Antiviral medicines (i.e. Tamiflu) can be prescribed to attempt to prevent symptoms of both types of influenza after an exposure.

BREAST-FEEDING WHILE ILL

Breast-feeding is strongly recommended for infants during the flu season. Infants who are not breast-feeding are more vulnerable to infection and hospitalization for severe respiratory illness than infants who are breast-feeding. If you become ill with influenza it will be very



important to continue to provide your baby with your breast milk either by nursing or by pumping to feed. While you are ill, it is best to have healthy family members or friends care for your baby. If you are ill and do not have anyone available to help with infant care, it is advisable for you to wear a mask, if possible, when caring for your baby. Antiviral medication (i.e. Tamiflu) is acceptable for use when breast-feeding. As always, contact your pediatrician for additional information.

ZIKA VIRUS – WHAT YOU NEED TO KNOW

- 1. Zika primarily spreads through infected mosquitos.
- 2. The best way to prevent Zika is to prevent mosquito bites.
- 3. Zika is linked to birth defects.
- 4. Pregnant women should not travel to areas with risk of Zika.
- 5. Returning travelers infected with Zika can spread the virus through mosquito bites and sex.

Zika virus updates and information can be found at www.cdc.gov/pregnancy/zika.

COVID-19

1. Know the facts:

- COVID-19 can spread between people who are in close contact with one another (within about 6 feet).
- Some people with COVID-19 may have no symptoms.
- Current reports suggest that pregnant women have a higher risk for more severe illness from COVID-19 than nonpregnant women.

2. Slow the spread:

- $\ ^{\blacksquare}$ Wear a mask or cloth face covering over your nose and mouth when in public.
- Clean hands often for at least 20 seconds with soap and water or hand sanitizer that contains at least 60 percent alcohol.
- Limit contact with other people as much as possible.
- Stay at least 6 feet away from other people if you need to go out.

Talk with your provider:

- Prenatal and postpartum care: Before an in-person visit, contact your provider if you think you may have COVID-19 or contact with someone who has it.
- Your birth plan: In most cases, the way you plan to give birth does not need to change. The safest place for you to give birth is still a hospital.
- Visitor policies: You may not be able to have as many visitors at your checkups or during and after birth while COVID-19 is spreading. For the most up to date visitor policy at the hospital, please call labor and delivery at 850-747-7700.

If you have tested positive for Covid-19, please call the office with the date your first symptoms appeared and the date of your positive Covid-19 test. Your scheduled appointment will be rescheduled to a later date. If your provider feels that you must be seen, we will give you appointment instructions at that time.

Herpes Infections & Pregnancy



Oral and genital herpes infections can be problematic in pregnancy. The biggest concern with genital herpes is transmission to the baby at the time of birth. Newborn herpes infection is rare but can be devastating to the baby. Transmission occurs by "shedding" of virus particles from an active herpes lesion. The risk of transmission is high if the mother has her first infection during the pregnancy (Primary infection). On rare occasions the mother can transmit a primary infection to her baby through the placenta during the first or early second trimester. This can cause serious fetal malformations and also lead to a miscarriage. Transmission to the infant with recurrent herpes infections is much rarer because the mother has usually passed antibodies to the baby through the placenta during the pregnancy.

Patients with an active genital herpes infection need to be delivered by c-section. If you have a history of herpes infection and present to the hospital in labor, it is important to notify your nurse and/or provider if you think you have symptoms of an outbreak. The virus is transmissible to the baby even in the first couple of days before a blister starts. It is recommended that patients with a history of herpes be started on an antiviral medication between 35-37 weeks gestation. Recent studies have shown that this significantly reduces the risk of an outbreak at the time of delivery. This medication is usually continued several weeks into the postpartum period.

About 5% of newborn herpes infections are contracted after birth. Make sure people always wash their hands before touching your baby, and never let anyone with a cold sore kiss, nuzzle, or hold your baby. Any herpes infection can be dangerous to a newborn, including the kind that causes cold sores. If you have an outbreak anywhere on your body cover it and minimize baby contact with that area. Wash your hands frequently. If you develop a herpes sore in or on your mouth avoid kissing your baby until it is completely gone. Wear a mask until the sore has crusted and dried.

It is safe to breast-feed while you have a herpes outbreak as long as there are no lesions on the breasts. Once again, hand washing in this case is imperative.



Headache in Pregnancy

Occasionally, a woman with chronic headaches prior to pregnancy will experience a reduction in headache frequency. However, due to increased estrogen levels during pregnancy some women who suffer from migraines may have an increase in the frequency and severity of their headaches. Unfortunately, most women with problem headaches will continue to live with headaches during their pregnancy. Due to hormonal changes, blood sugar fluctuations, and sometimes, dehydration, women with <u>no</u> history of migraines may develop headaches during pregnancy.

TIPS FOR LIVING WITH HEADACHES

- If you are under the care of a neurologist, chiropractor, or pain management specialist, please continue your care while pregnant.
- If you need medication for your headaches be sure to talk with us about it.
 Some headache medications should not be used during certain periods of your pregnancy.
- Avoid headache triggers. Some food triggers include; MSG, chocolate, nuts, peanut butter, bananas, onions, avocado, NutraSweet (Equal), pickled foods, yogurt, sour cream, aged cheese, processed meats, pizza, and caffeine.
- Eat frequent small meals that contain some protein. Examples of protein include dairy products, eggs, meats, and beans. These foods help you avoid low blood sugar which is a common cause of headaches during pregnancy.
- Drink plenty of liquids. Dehydration is a common problem during pregnancy and can contribute to headaches. Water is your best source of hydration. Remember that small amounts of caffeine have not been shown to be harmful during pregnancy; however, it is a diuretic and can cause dehydration. Furthermore, it is not uncommon for patients to experience headaches secondary to caffeine withdrawal.
- 400 mg of Magnesium Oxide taken twice daily, has been shown to significantly decrease incident of migraine headaches.
- If you think your headache may be related to sinus congestion try using a decongestant (see page 24). You can also place warm compresses on your face just above and below the eyes. Saline nasal sprays and breathing steam can also help reduce the pressure of sinus congestion.
- It is OK to take Tylenol alone or with Benadryl (Tylenol PM) to treat your headache. Do not take Benadryl or Tylenol PM unless you can rest for a while in a darkened and quiet room. Benadryl causes sleepiness and is unsafe to use if you are doing anything that requires you to remain alert. If you need to remain alert, take plain Tylenol regular or extra strength as directed on the bottle.


***If you are having a severe headache accompanied by visual changes, weakness, or numbness, contact us immediately even if it occurs after your delivery.

Ligament Pain



There are 3 main sets of ligaments that support the uterus. 1) The round ligaments attach the front sides of the uterus to the pubic bone. 2) The broad ligaments support the far sides of the uterus to the pelvic bone sidewalls. 3) The uterosacral ligaments support the back of the uterus to the lower part of the vertebrae just above the buttocks. As the uterus begins to grow out of the pelvis, the ligaments stretch like thick rubber bands and can cause cramping and sharp pains.

- During the first three months of pregnancy many women report that they feel crampy just like they do before or during a menstrual period. This may be due to the stretching of all 6 of the ligaments.
- During the second and third trimester the round ligaments can cause pain in the lower abdomen, in the pubic region, and on either or both sides. This is a normal but uncomfortable occurrence and usually resolves once the uterus grows to full-term size. When round ligament pain occurs, there is a sharp, stabbing kind of pain, that fortunately is brief. Round ligament pain frequently occurs with a sudden movement, coughing or sneezing, stretching, heavy lifting, standing, or a twisting of the upper body (like beginning to roll over in bed without first bending the knees.) When round ligament pain occurs it can be resolved by lying on the side that hurts - with the legs curled up against the abdomen. A heating pad, warm bath, and/or Tylenol frequently help relieve the discomfort. An elastic maternity belt is also helpful and can be worn over or under your clothes. If the pain is prolonged without improvement, or is accompanied by vaginal bleeding, fever, upper/mid back pain, nausea, vomiting, diarrhea, or burning with urination you should contact us or report to Labor and Delivery.
- Low back pain is usually caused from stretching of the <u>uterosacral ligaments</u>. This can frequently be eased with a heating pad, warm bath (remember that prolonged soaking in hot tubs is not recommended during pregnancy), massage, and Tylenol. We also recommend the "Cat & Cow" Yoga exercise to relieve this discomfort. Please review recommendations found in the back pain section of this book on page 43.



Back Pain During Pregnancy



Back pain is a very common complaint during pregnancy. As your baby grows there will be strain and stretching of the back muscles, ligaments, and joints. Additionally, hormonal changes of pregnancy will cause some relaxation of your abdominal muscles. Furthermore, pregnancy changes your center of gravity and causes you to walk differently making you use muscles you are not used to using.

To help prevent or ease low back pain, try our suggestions listed below:

- Be aware of how you sit, stand and move. Check your posture. Tuck your tummy in and keep your back straight, as if your hair is being pulled straight up.
- Wear low heeled shoes with good arch support.
- When standing for long periods, move in place or put one foot up on a stool.
- If your bed is soft, have someone place a board between the box spring and mattress.
- Try to not bend from the waist to pick things up, squat down, bend your knees and keep your back straight.
- Sit in chairs that have good back support. Try placing a small pillow behind your low back when sitting.



- Sleep on your side with pillows between your legs for support. Try a small pillow under your tummy.
- A hot or cold pack applied to the area may help. Try a warm shower or bath
- Book a massage, or get your partner or a friend to help. If you want to book a massage, most licensed therapists require a prescription before they will work on pregnant patients, so let us know and we will refer you.
- Use a maternity support belt to help support the growing uterus and your low back muscles.
- Do the "Cat & Cow" Stretch exercise for low back pain
 - ▶ Get on your hands and knees and curl your back up like an angry cat, holding the position for the count of 5 while blowing out all of your air.
 - ► Then, flex your back down toward the floor letting your back sway like a cow. Look up to the ceiling and breathe in deeply.
 - ▶ Breathe in when flexing to the floor and breathe out when arching up.
 - ▶ Let your baby dangle and concentrate on relaxing your spine muscles.
 - ▶ Repeat these stretches 15 to 20 times at each session and do them several times a day.

Cat & Cow Stretch





- Do the Pigeon Stretch exercise for sciatic and low back discomfort
 - ▶ Start on your hands and knees with your hands shoulder length apart.
 - ▶ Bring your left leg in front of you with the knee bent. Bring the left knee to the outside of the left and allow the right leg to straighten and release to the floor.
 - ► Square the hips towards the floor.
 - ▶ Add padding under the left side of the buttock as necessary to bring the hips square.
 - ➤ You can then extend your right foot backward and deepen the stretch while keeping your hips straight.
 - ▶ Hold the stretch for several minutes then switch sides.

There is a link to a short instructional video in the FAQ section of our website. www.emeraldcoastobgyn.com

Remember, sometimes back pain can be the first sign of a serious problem. If you find that these measures are not helpful for your back discomfort or if you are becoming worse rather than improving please call to discuss your symptoms.




Exercise During Pregnancy

In the absence of medical or pregnancy complications, 30 minutes or more of moderate exercise a day is recommended. In general, pregnant women engaging in regular, moderately intense physical activity continue to derive the same health benefits during pregnancy as they did prior to becoming pregnant. Some women are very physically active and are used to regular exercise. With a few modifications these women may continue their program as long as their pregnancy is uncomplicated. Women who are not used to regular exercise but want to begin an appropriate program may do so to gain the health benefits that exercise can offer during pregnancy. Exercise increases your energy level, helps mobilize fluids to decrease swelling, and may be helpful in the prevention and management of gestational diabetes. As always, if you have any questions or concerns regarding a specific exercise program or any problems you encounter during exercise, please talk with us.

CONSIDERATIONS AND GUIDELINES

- Remember that your center of gravity changes as the pregnancy advances which can make you clumsier and more prone to falling. Plan accordingly and try to exercise with a buddy or at least carry a cellphone.
- Whenever you exercise remember that you need to increase your fluid intake so that you avoid dehydration, dizziness and muscle cramps (remember, that the uterus is a muscle!).
- Be sure to wear shoes and clothing appropriate for your exercise routine.
- Stop the activity, or slow down and rest if you become extremely short of breath, develop pain in your chest, notice dizziness, or develop frequent uterine contractions. If these symptoms continue, speak with us about the problem.
- Many women find that walking, dancing, swimming, and Yoga offer excellent benefits and are ideal exercises during pregnancy and the postpartum period. After the first trimester, try not to exercise while lying flat on your back except for brief moments. We recommend the Gaiam Prenatal Yoga app called "Yoga Studio". With this app you can enjoy a free trial for 7 days and they offer monthly and yearly subscriptions.
- Avoid sports with a high potential for contact injuries such as basketball, football, and soccer. Also avoid exercise with a high potential for falling such as horseback riding, mountain climbing, gymnastics, downhill skiing, and cycling. This is especially important in the second half of pregnancy when the baby loses the protection of the pelvic bones.

- Scuba diving below 32 feet during pregnancy is not recommended. Your fetus can get the bends even if you do not. Skin diving and snorkeling are fine, but remember to use the buddy system. You will fatigue easier during pregnancy.
- Avoid bouncy or bumpy rides such as jet skiing or boating in choppy water, and off-road riding in all-terrain vehicles.
- When participating in aerobic exercise be certain to slow down whenever you notice that you cannot talk comfortably during your workout.
- Always wear protective clothing and sun screen while exercising outdoors.





Travel During Pregnancy

Normal pregnancy should not put a damper on travel plans but some precautions are necessary. The best time to travel is between weeks 14 and 32. During this time you are usually less queasy and are not as likely to experience some of the common discomforts of late pregnancy. If you do plan to travel, it is recommended that you carry a copy of your prenatal records with you in the event you need to be seen in a hospital during your trip. Please arrange to see us shortly before you leave and a copy of your records will be given to you on request.

If you are traveling by plane, most airlines will allow you to fly up to a month before your due date. Try to get an aisle seat towards the front of the plane. Don't worry about the metal detectors at the airport. They will not harm your baby. If you are traveling to a foreign country please discuss this with your provider. Women who are pregnant should not travel to areas with active Zika virus infections reported. Please go to http://www.cdc.gov/pregnancy/zika/ for the most up to date information regarding the Zika virus.

TRAVEL TIPS

- Always wear your seatbelt snugly across your pubic bone and below your growing uterus when in an automobile. Do not disable your air-bags.
- Before your trip, ask your family or friends for the name & phone number of an obstetric provider, or a hospital with obstetric services.
- Keep your travel plans flexible. Problems can come up at any time. Buy travel insurance to cover tickets and deposits that can't be refunded.
- While traveling, plan to walk about every hour. Stretch your legs often to
 lessen the risk of developing a blood clot and make you more comfortable.
 This will also decrease the amount of swelling in your ankles and feet. Don't
 cross your legs while sitting. Don't sit in the same position for long periods
 of time.
- If you are traveling longer than 4 hours, one 'baby aspirin' (81mg) taken the day before your trip and on the day of travel may help prevent blood clots.
- Wear comfortable clothing & shoes. Consider wearing support stockings.
- Carry light snacks and water with you.
- Take time to eat. A balanced and healthy diet during your trip will boost your energy and keep you feeling good. Be sure to get plenty of fiber to ease constipation, a common travel problem.
- Drink plenty of fluids to help prevent dehydration, uterine cramping, and urinary tract infection. Empty your bladder at least every two hours.
- Get plenty of sleep, and rest often.
- Stretch your back and neck muscles from time to time.
- Don't do too much. It's tempting to squeeze in as many sights as you can, but it is important to adjust your pace when you are traveling during pregnancy.

Fetal Kick Counts



Most pregnant women begin to feel the baby move sometime between the 16th and 20th weeks of pregnancy. Women who have had babies in the past will frequently feel this movement earlier than those having their first baby. We call this early movement "Quickening", a term that comes from the Bible - meaning "life". Babies are usually the most active between the hours of 10:00 PM and 2:00 AM. They may sleep for up to 40 minutes with the sleep cycle lengthening as the pregnancy progresses. Also, during late pregnancy your baby may decrease the big kicks and huge flips simply because of running out of playground space. Movements may become softer but should continue to be frequent and reassuring. After the 24th week of pregnancy keeping track of your baby's movement habits becomes an increasingly important way for you to monitor your baby's health. If, after the 24th week of your pregnancy you feel that your baby is not moving normally, please do a *Fetal Kick Count*. Simply follow the guidelines listed below.


FETAL KICK COUNT GUIDELINES

- Before beginning, be certain that you have recently nourished yourself. If not, eat a healthy snack with a glass of water, milk or juice.
- In a quiet place, lie down on your side or sit in a comfortable chair. Avoid the distractions of work, television, conversations, and interruptions.
- Pay close attention to the movements of your baby. Check the time. Count every kick or movement until your baby has moved at least 10 times. Healthy babies can sleep for up to 40 minutes out of an hour! For this reason, it is important to give your baby up to 2 hours to complete the test. Once you count 10 fetal movements you can go about your day with reassurance. Remember that your baby's sleep cycles may lengthen as your pregnancy progresses.
- As your baby grows larger, the movements usually become softer and gentler. All of your baby's movements count, even if the activity is not as prominent as you may have noticed before.
- If you have completed 1-4, above, and your baby has still not moved at least 10 times within 2 hours you should come to our office for electronic fetal monitoring. If we are closed, you should go to Labor & Delivery at Gulf Coast Hospital. Call us first so we can let the nursing staff know you are coming. This will give us reassurance that all is well with your baby.



Urinary Tract Infection

Urinary tract infections can be serious during pregnancy. Bodily are taking place increase your risk of urinary tract Pregnancy hormones infection. relax the muscles and valves in the urinary system, which can cause urine to move more slowly through the urinary tract. Furthermore. pressure from the growing uterus can squeeze or pinch the small tubes (ureters) that carry urine from the kidneys to the bladder. Bacteria. once in the bladder can travel upwards toward the kidneys, and can cause a very serious infection. In addition to making you very ill, a kidney infection can trigger pre-term



labor. Pregnant women with kidney infection usually require hospitalization with intravenous antibiotics, as well as low dose antibiotic throughout the remainder of the pregnancy to prevent recurrence.

- SYMPTOMS OF A BLADDER INFECTION: Pain and/or burning before, during, or after urinating. Feeling more than the usual urge to urinate or increased frequency of urination. Sometimes blood can be seen in the urine or on the toilet tissue. The urine may or may not have a foul odor. Please report these symptoms to us.
- SYMPTOMS OF A KIDNEY INFECTION: Frequently the symptoms are the same as for a bladder infection but are often accompanied by fever, chills, generalized aches and pains, and pain in the mid-back or side. Women with a kidney infection sometimes feel like they are coming down with the flu.
- SYMPTOMS OF A KIDNEY STONE: This can occur with or without an infection. The most frequent symptoms are severe colic-like or spasmodic pain that does not improve with movement or time. The pain can be located in the back, side, lower abdomen, or the pubic area. Sometimes there is bloody urine or blood on toilet tissue. Please report these symptoms to us.

HOW YOU CAN HELP PREVENT URINARY TRACT INFECTIONS?

- Always wipe from front-to-back instead of wiping germs from the anal and vaginal region toward the urethral opening.
- Drink plenty of liquids daily so that you need to empty your bladder about every 2 hours. Avoid caffeinated liquids because they can cause dehydration. Cranberry juice has been reported to reduce your risk of a bladder infection.
- Always empty your This helps flush t urethra of germs.
- When lying down, remember to lie on your side. This will help release pressure off of the kidneys and the ureters.
- Rest in the knee chest position, especially if your mid-back region is hurting.
- Make sure you are getting plenty of vitamin C.
- If antibiotics are prescribed, be certa to take all of thei and take them at th times prescribed Never quit taking the antibiotics because you feel better! The surviving bacter will become resist to that antibiotic.





Pre-term Labor

Pre-term labor is defined as the onset of consistent uterine contractions that are every 5-8 minutes in a woman who is between 20-36 weeks pregnant, who in addition, has either spontaneous rupture of the membranes or progressive cervical dilation. Most women who have pre-term contractions do not go into pre-term labor.

Premature birth occurs in about one of every ten pregnancies. Prematurity can cause health problems for the baby because their lungs and other organs have not had time to fully develop. The earlier a baby is born, the more likely he or she will have health problems related to pre-term birth. Some of these problems can be immediately life-threatening as well as cause problems that affect the infant throughout his or her life.

WOMEN ARE AT INCREASED RISK OF DELIVERING PRE-TERM IF THEY:

- Have delivered a premature baby in the past.
- Are habitual users of alcohol, tobacco, or street drugs.
- Have had infections that are not treated.
- Have a multiple gestation pregnancy (i.e. twins.).
- Weighed less than 100 pounds before becoming pregnant.
- Are younger than 18 or older than 40.
- Have poor nutrition, poor dental hygiene, or obesity.
- Had bleeding problems during the pregnancy.

NOTE: Just because you don't have any of the above risk factors DOES NOT mean you can't have pre-term labor.

PRE-TERM LABOR SIGNS AND SYMPTOMS

- Irregular or regular contractions (like menstrual cramps).
- Low backache.
- Increased vaginal discharge, spotting, or fluid leaking from the vagina.
- Feelings of pressure in the pelvis.
- Fever (oral temp greater than 100.5)

IF YOUR SYMPTOMS ARE MILD AND YOU HAVE NO FLUID LEAKING OR BLEEDING YOU MAY TRY THE FOLLOWING:

- Stop what you are doing and find a place to relax.
- Drink two or three glasses of caffeine free liquid (water, juice, Gatorade).

- Empty your bladder.
- Sit or lie down and take note of the frequency of your contractions for one hour.

If your symptoms go away, you can go about your regular activities but continue to monitor for contractions. If the symptoms do not go away and you notice more than 4 painful contractions per hour call the office. During business hours we may have you come to the office for evaluation. If the office is closed we will direct you to Labor & Delivery. Once again it is very important for you to call our office (850-769-0338) and let the on-call provider know you are on your way.

- During an evaluation of pre-term labor we will monitor your contractions using the electronic fetal monitor and may perform a vaginal ultrasound to measure the length of your cervix. A short cervix can be a risk factor for pre-term birth. We sometimes also order a test call Fetal fibronectin to determine your risk of delivering prematurely within the next few weeks. Each of these tests is a non-invasive method of evaluating your risk for premature labor. These tests are helpful but not definitive. They frequently will give false positive results. Pre-term labor has many causes and often can be a difficult problem to diagnose.
- ▶ Patients who have a history of pre-term birth are at increased risk of pre-term labor in subsequent pregnancies. Weekly progesterone injections starting at 16-18 weeks have been shown to decrease the risk of pre-term delivery in this high-risk group of patients. If you feel you may be at risk for pre-term labor, please discuss this with your provider.





Blood Pressure Problems

Blood pressure (BP) refers to the amount of pressure exerted on the walls of your arteries during both the contraction and relaxation phases of your heart-beat. A blood pressure reading has 2 numbers. You may hear it referred to as "120 over 80". The first number is called the *systolic* pressure, and the second number is the *diastolic* pressure. Blood pressure changes often throughout the day. When a person's blood pressure stays high for some reason, it may signal a problem.

If you are pregnant and your BP is 140/90 on several readings, this is too high. When a pregnant patient has high blood pressure it can cause blood flow problems to the fetus which can compromise growth or cause other problems with the placenta.

CATEGORIES OF HIGH BLOOD PRESSURE

CHRONIC HYPERTENSION:

This condition occurs when a mother is diagnosed with high blood pressure before pregnancy. Often medications have to be changed to drugs that are safer in pregnancy. Due to cardiovascular changes throughout the pregnancy, patients with chronic hypertension have to be monitored very closely.

GESTATIONAL HYPERTENSION:

When patients who do not have chronic hypertension, develop high blood pressure during the second half of pregnancy it is referred to as gestational hypertension. This condition requires careful monitoring and can lead to pre-eclampsia (see below).

PRE-ECLAMPSIA (previously referred to as Toxemia)

This is a serious medical condition that can affect all organs of the body. We do not completely understand why pre-eclampsia occurs, but it is a potentially life threatening disease that occurs in the presence of high blood pressure usually in the third trimester. It is usually diagnosed by the presence of protein in the urine representing stress on the kidneys. Other clinical features of pre-eclampsia include:

- Headaches
- Rapid weight gain

- Severe swelling of hands and face
- Visual disturbances
- Abdominal pain around the liver
- Seizures if seizures develop, the disease is referred to as Eclampsia

Patients with pre-eclampsia may need to be hospitalized. Patients that develop severe pre-eclampsia may require early delivery.

If you develop any blood pressure problems in pregnancy, you can expect more frequent office visits and increased antenatal surveillance (see the section of the booklet regarding fetal testing).



Gestational Diabetes Mellitus

Diabetes is a disease that causes high blood sugar. Diabetes can occur in both pregnant and non-pregnant women. When it occurs in pregnancy, it is called gestational diabetes and if a patient is diabetic before pregnancy it is referred to as pre-gestational diabetes. There are serious health problems and complications that are associated with diabetes in pregnancy. Gestational diabetes usually resolves with the birth of the baby; however, it may persist or return later in life. This is more likely to occur in women who do not maintain a healthy diet and normal body weight as they mature.

To screen for diabetes your provider will order a 1 hour glucose tolerance test (GTT). This is usually ordered between 24 and 28 weeks. Your provider may decide to do an "early glucose tolerance test" if you have certain risk factors. If you have an early GTT, you will still need the regularly scheduled GTT in the second trimester. If your 1 hour GTT is abnormal, you will need to undergo a 3 hour GTT to make a definite diagnosis of diabetes. You do not need to be fasting for the 1 hour test but follow our eating directions found in the laboratory testing section on page 19. For the 3 hour GTT, you must be fasting! Nothing to eat or drink after midnight except water until your test is completed. We encourage water. Not drinking enough water makes multiple blood draws difficult. Otherwise, no gum, no anything!

The ultimate goal in the treatment of pregnancy associated diabetes is overall control of the mother's

blood sugar. Optimal blood sugar control has been shown to significantly reduce the risk of inherent pregnancy and delivery complications.

After your baby is born, you will need to be tested again. This is to screen for non-gestational diabetes



If you are diagnosed with gestational diabetes, you will be scheduled for out Gestational Diabetes Class, taught by a Master of Public Health, Registered and Licensed Dietitian Nutritionist. Emerald Coast OB GYN is the only obstetric office in the area to offer this type of specialized class. Be sure to attend if you are scheduled to help you learn how to how to manage your gestational diabetes with healthy food choices.

Abdominal Trauma



Pregnancy provides no immunity against accidents and injury. In fact, pregnant women are more prone to falls due to changes in the center of gravity and weight gain. It is natural to worry more about any injury sustained during pregnancy because of fear of harm to your fetus. Most injuries do not damage the baby or cause the onset of miscarriage or labor. However, there are particular types of injuries or accidents that do cause us concern and must be carefully evaluated. Often the concern is related to the stage in pregnancy which the injury occurs. For example, during the first 12 weeks of your pregnancy, the developing embryo is quite well protected due to the thickness of the uterine muscle and the structure of your pelvic bones. As your pregnancy develops, the uterus becomes an abdominal organ and the uterine muscle becomes progressively thinner. Therefore, the farther along you are, the more serious the injury can be, especially when abdominal trauma and falls are involved. Here is our advice:

FIRST TRIMESTER:

If you have a motor vehicle accident or take a fall during the first 12 weeks of pregnancy you will most likely not have injured your baby. Remember that during this period of time it is normal to experience crampy pelvic and low back pain similar to those many women have with their menstrual periods. If you end up in the Emergency Room, they will check your fetal heart tones before they release you. If you have a minor injury and don't need to go to the Emergency Room, use common first aid measures such as 'RICE', (Rest, Ice, Compression, and Elevation). You may take Tylenol for minor aches and pains. Notify us if you experience vaginal bleeding.

SECOND AND THIRD TRIMESTERS

(After 13 weeks gestation):

If you have a significant fall, abdominal trauma, or an auto accident that is more than a fender-bender (i.e. you were traveling at more than 20mph, your air bags deployed, or you were compressed against the steering column or dash), we will want to evaluate both you and your baby at GCMC. While most severe obstetric complications are diagnosed within the first 4 hours following abdominal trauma, some can develop over a longer period of time. For this reason we may want to evaluate you for several hours immediately after the injury. If you are out of town at the time of the accident or injury, you should report to the nearest hospital that has an obstetric department. If you sustained a serious injury to another part of your body, such as a head or neck injury, severe laceration, fracture, or other significant bodily injury, you should receive a triage assessment in the Emergency Room prior to being released to Labor & Delivery for evaluation of your baby.



Additional Fetal Surveillance

Sometimes we need to increase the level of fetal wellbeing assessments over-and-above routine monitoring with fetal kick counts by additional testing. These tests are non-invasive in nature and are used to detect problems that may arise. There are many reasons why your provider might order additional fetal surveillance. The following list contains examples of some reasons these tests might be ordered.

PRE-EXISTING MATERNAL PROBLEMS

- Chronic hypertensive disorders
- Severe heart disease
- Chronic renal disease
- Poorly controlled hyperthyroidism
- Maternal age above 40 years

- Clotting disorders
- Type 1 diabetes mellitus
- Lupus and other systemic diseases
- Hemoglobinopathies, such as Sickle Cell Anemia

PREGNANCY RELATED DISORDERS

- Pregnancy induced hypertensive disorders
- •Too much or too little amniotic fluid Fetal growth problems
- Gestational diabetes
- Previous stillbirth
- Overdue baby

- Decreased fetal movement (see our page on fetal kick counts)
- Multiple gestation
- Placental problems
- Blood or immune disorders

TYPES OF TESTING

NON-STRESS TEST (NST)

This is an assessment of a baby's well-being. By changes in the fetal heart rate pattern, your provider can verify with a large degree of certainty that your baby is OK.

AMNIOTIC FLUID INDEX (AFI)

This is an ultrasound test to measure the amount of fluid around your baby.

BIOPHYSICAL PROFILE (BPP)

This is an ultrasound test to evaluate the baby's movement, muscle tone, breathing motion, and amniotic fluid volume. This test helps us determine the baby's overall well-being.

Group B Strep Testing



Group B Streptococcus (GBS) is a common bacterium that lives within the human body without causing harm to healthy people. GBS can be found in the intestine and vagina in about 3 of every 10 women. It is not a sexually transmitted disease, and it does not usually cause discharge, itching, odor, or other symptoms. It will not harm you or your sexual partner. However, in rare circumstances GBS can seriously harm a newborn baby. At the time of birth, babies are exposed to the GBS bacteria if it is present in the vagina. This can result in pneumonia, meningitis, or a blood infection. Full-term babies who are born to mothers who carry GBS in the vagina at the time of labor and childbirth have a 1 in 200 chances of getting sick from GBS during the first few days of life. Premature babies have an increased chance of becoming ill from GBS if it is present.

We can discover most people at risk for GBS transmission by culturing every pregnant woman shortly before the expected due date. We recommend culturing between the 35th and 37th week of pregnancy or earlier is there is a risk of pre-term labor (twins, etc.) For those patients anticipating a Cesarean section, we still advise the GBS test because this information will be helpful in treating any baby who becomes ill - regardless of the route of birth. We obtain the culture at the office during a regular prenatal visit. Your results will be available to us and to the nurses in Labor & Delivery after 48 hours. Occasionally, GBS can cause a urinary tract infection during pregnancy. GBS urinary tract infections are treated at the time of diagnosis.

If your culture is positive, we will recommend that you receive antibiotics during labor. If you carry GBS at the time of birth and you are given intravenous antibiotics during labor, the risk of your baby getting sick from GBS is 1 in 4000. If a baby becomes sick with GBS, it usually occurs within the first 24 hours after birth.

If you go into labor before you are cultured for GBS we will consider your GBS status 'unknown' and will recommend that you receive antibiotics during labor. In some circumstances, GBS positive patients may not have adequate time to get sufficient doses of antibiotics. In these cases, your baby may be admitted to the NICU for an observational workup for infection.



Vaginal Birth After Cesarean (VBAC)

If you have had a C-section in the past, it is imperative that you discuss the topic of VBAC with your provider. This term refers to the process of giving birth vaginally when you have previously had a Cesarean delivery in the past. Approximately 70% of women who attempt a VBAC will successfully deliver vaginally. The remainder will ultimately require a repeat C-section. Depending on your specific medical condition, there may be additional risks, benefits, and/or alternatives to VBAC. It is important to discuss your individual risks, options, and concerns with your clinician. Women with a history of having a vaginal birth prior to their previous cesarean are better candidates for VBAC.

Benefits of VBAC:

VBAC has a smaller risk of complications such as hemorrhage and infection, for the mother, when compared to repeat C-section. Recovery time is shorter and usually associated with less pain.

Risks of VBAC:

The greatest risk associated with VBAC is the increased risk of uterine scar separation, or rupture, at the site of the previous Cesarean incision. Such a separation may be minor or catastrophic requiring emergency Cesarean delivery. Most studies report the risk of uterine rupture to between 1%-3%. More recent studies place this risk as low as 0.3%-0.4% if the previous cesarean was not a vertical incision on the uterus. Emergency Cesarean delivery becomes a lifesaving procedure in cases of uterine rupture. The emergency nature of the operation adds greater risk to the mother and baby more than a planned repeat Cesarean birth.

VBAC is associated with a higher risk of harm to the baby than to the mother. If the uterus ruptures during a VBAC attempt, there may not be enough time to operate to prevent death or permanent brain injury to the baby.

Consequences to the woman who experiences a uterine rupture can include hemorrhage, damage to the uterus, bladder, ureters, bowel, and possibly death. In such an emergency there is an increased risk for the need for a blood transfusion and the risk of post-operative infection is also higher.

Selecting a Pediatrician



Choosing a pediatrician is generally a matter of personal preference and insurance restrictions. Like providers in other specialties, pediatricians may participate in some insurance plans but not others. We are very fortunate here in Panama City to have many skillful, kind and respected pediatricians in practice. Check the telephone book's Yellow Pages or ask other family members and friends for their recommendations. In addition, you may call the Gulf Coast Medical Center's, "Consult-a-Nurse Health Care Referral Line" at 747-3600 for suggestions and answers to common questions that you may have about locating a pediatrician in our community.

If you know which pediatrician you want to have care for your baby, you should call their office by your 32nd week of pregnancy to make sure they will be able to accept your new baby into their practice. You may want to request an office visit to sit down and talk about office policy and other subjects important to you - so that you get a chance to become acquainted before your baby is born.

Rather than make this type of advance arrangement, many expectant parents choose to use the services of the "on-call" pediatrician for their baby's immediate newborn care. All pediatricians practicing at Gulf Coast Medical Center take turns being the "on-call" pediatrician and provide care to all babies born during that time. Most parents find they like the "on-call" pediatrician so much that they continue care in that pediatrician's office long after their newborn baby is discharged from the hospital's nursery. Tricare families find that this option seems to work out for the best because most military families will transfer care to a physician on-base for follow-up pediatric care.



*We would like to thank the pediatric offices who have helped sponsor this educational booklet by placing an advertisement in the back of our book. We appreciate their commitment to patient education!



Induction of Labor

Inducing labor means starting or bringing on labor through medications or other methods before labor begins on its own. Labor may be induced for a variety of reasons. A primary risk of inducing labor is that it may be unsuccessful. If an induction is unsuccessful, a Cesarean section becomes necessary. When compared to a vaginal birth, a Cesarean carries higher risks of postpartum complications and a significantly longer recovery period. Generally, inductions are separated into those that are medically necessary and those that are elective.

MEDICALLY INDICATED INDUCTION:

This type of induction is recommended when the health of the woman or baby is at risk. The following are examples of why a medical induction might be needed: Overdue baby, blood pressure problems, infection, placental problem, ruptured membranes. Sometimes, an induction may be recommended due to concern that you may not make it to the hospital in time once you go into labor.

ELECTIVE INDUCTION:

This type of induction is one chosen by the woman for convenience or other non-medical reasons. It is important to understand that if you opt for elective induction, there are some associated risks. This is an important decision that you should discuss extensively with your doctor or midwife. Ideally, elective inductions are scheduled at about 39 – 41 weeks of pregnancy (by our best estimate) and when the cervix is determined to be favorable. We can evaluate your cervix when performing a pelvic exam during late pregnancy. Induction initiated when the cervix is not 'favorable' or 'ripe' carries an increased risk of the need for a cesarean section. In order for you to be a candidate for elective induction, your provider must be sure of your due date. Patients who present for initiation of prenatal care late in pregnancy often are not ideal candidates for elective induction.

Inducing labor before the cervix is ready requires use of cervical ripening medication and necessitates hospital admission for the ripening - before the actual induction can begin. This adds expense to the childbirth process. Perhaps the best way to be certain that your baby is ready to live outside your body is to wait for the baby to initiate its own labor (within a reasonable time-frame). Inducing labor too early can place the baby at increased risk for immaturity problems such as breathing difficulty, temperature control problems, newborn jaundice, and problems coordinating suck and swallow.

These problems can cause separation if your baby needs to remain in the nursery or the NICU for close observation. It can also complicate the initiation of breast-feeding, and sometimes contributes to a longer hospitalization for baby or readmission to treat for jaundice.

INDUCTION SCHEDULING

When we schedule a labor induction, there is no guarantee that Labor & Delivery will have space for you! Unlike reserving a hotel room, the Labor & Delivery unit cannot "hold" a room for us. Women who present in natural labor or require admission for medical necessity will have first priority, just as you would in the same situation. Sometimes elective inductions get "bumped"! This can be frustrating for us and especially for you and your family. For this reason, it is important that the Labor & Delivery staff has a working emergency telephone number where you can be reached if changes become necessary. Also, it is recommended that you call Labor & Delivery to be certain a room is available before you drive to the hospital. The phone number for Labor & Delivery is 850-747-7700. If your induction has been postponed, contact our office the next morning after 8:00

Questions: Please ask us any questions you may have about possible induction of labor. We consider the advisability of induction on an individual basis and will be happy to answer any questions you may have in this regard.

e fmsx brndsecpeog c i rdrmipfloecks e s s c a t t l r u n u t e l t u tudobstetricians s m i c m a y r o h t d b t u a d e l e s o a e dan r i i o e om arb cyglueeupcm e b t v t d a i o a m s m s k a wriii lgssgeueocg aaoxnaattnctehir c n n s m r f e d u a t u r k c ue set enrsrfhlbbm t s n l e m r p d e f i w d i m wt udielyteeeyens s u r e t u hua f m d b s l l bu ixg esfnund coab

ultrasound emeraldcoast hospital kickcounts cervix uterus girl boy ligaments mucusplug membranes fetus hydrate contraction midwife breastfeed birth dilation effacement preregister obstetrician

Across

1. Drink lots of these to stay healthy!

3. What usually lasts 280 days?

6. Our "Phone____" can answer most questions.

7. We use ____ Coast hospital.

schedulers can schedule your visits.

Down

2. The lab we prefer.


3. ____vitamins are important to take.

4. www.ecobgyn.com is our address

5. Pregnant women need their shots!

8. Your unborn baby is called a _____.

9. We recommend the South Beach



Broken Water?



About 10% of the time your water will leak or break before labor begins. If your water breaks (rupture of membranes), labor usually begins soon after. Usually rupture of membranes is obvious, however, on occasion the signs can be subtle. Sometimes, an examination is needed for confirmation. Here are some tips to help you decide what to do if you suspect your water has broken or is leaking.

- The first thing to remember is that leaking urine can seem like leaking amniotic fluid. The baby can head-butt your bladder and cause some urine to leak out.
- Importantly, when your bag of water is leaking it will **not** stop. If it is urine leaking, it will stop once you have completely emptied your bladder.
- Before coming to the office or to Labor & Delivery you can always take a few minutes to do the "Panty Test" to help you decide.

THE PANTY TEST

- Completely empty your bladder.
- Cleanse the vaginal region with soapy water, rinse off and dry completely with a clean towel (you may want to use a hair dryer to dry all your pubic hair completely).
- Put on the clean panties or a fresh panty liner and walk about your home for about 15 minutes.
- Return to the bathroom and check your panties or liner.
- If you remain wet after taking the above steps you will need to report to the office or Labor & Delivery for more specific testing.
- If you remain dry, go about your regular activities but keep alert for continuing signs of leakage. Sometimes the symptoms may be subtle. If you are ever in doubt, err on the side of caution and see your care provider.





True Labor?

Our recommendations outlined in this section are specific to labor at term. If you are experiencing these symptoms before your 36th week of pregnancy, please refer to our section on "Pre-term Labor". (pg. 53) If you are still indoubt, please give us a call.

Sometimes it can be difficult to determine if the symptoms you are experiencing are caused by 'pre-labor' or the real thing! Here are a few points of advice. If ever in-doubt you may certainly call us at 850-769-0338 or report to Labor & Delivery for a complete assessment.

PRE-LABOR SYMPTOMS

The nature of pre-labor or "false labor" symptoms are that they generally come and go. It can be uncomfortable but you will not see a progression in strength, intensity, regularity, or frequency over a period of time. Your contractions will usually be somewhat irregular in timing and strength – no real or predictable pattern. Another clue is that pre-labor or false labor contractions change in nature when you change your activity level. Test it by resting or walking. True labor will continue despite any actions you may take, whereas pre-labor or false labor symptoms will usually subside with activity change and hydration.

THE WATER TEST

To help you rule-out pre-labor or false labor, you can try this test. Drink 3-4 glasses of clear liquid and lie on your side for about one hour. If you are in pre-labor or false labor, your contractions will begin to fade or taper off. If you are in true labor, no amount of liquids or rest will change your symptoms or stop your contractions! As always, if you are unsure, you may always call us or report to Labor & Delivery. If you go to Labor & Delivery for assessment and end up being sent home, do not be embarrassed or angry! This happens frequently. Sometimes differentiating labor from false labor can be difficult, especially for first time mothers.

TRUE LABOR

The definition of true labor is regular and painful uterine contractions that cause the cervix to dilate - over time. The nature of true labor is that it is not influenced by anything you do. It will not matter if you remain still or active – true labor will continue. Real labor contractions may begin as mild

or irregular but over time become predictable, organized, and increasingly uncomfortable. If your water remains intact, time the contractions for at least one hour before deciding to come to the hospital. We want you to come to Labor & Delivery once your contractions are every 5 minutes and progressively getting stronger. As always, if your water breaks you need to go to the hospital without delay.

5-1-1

Remember the 5-1-1 rule! When contractions are no more than 5 minutes apart, lasting for at least one minute for a duration of one hour – you should start making arrangements to get to the hospital.

If you are in early labor and your water has not broken, you may want to walk, shop, take in a movie, catch up on household chores, or do some other activity to help you relax. A warm bath might be your perfect choice!

THE EXCEPTIONS

If you have a history of rapid labor (less than 3 hours), if you live more than one hour from the hospital, or if your Group B Strep Culture was positive, you will want to leave for the hospital once contractions are between 5 and 10 minutes apart. Mild bloody discharge can be normal up to two weeks before -- or with the onset of labor. This is sometimes referred to as "passing the mucus plug". However, if you have bleeding enough to saturate a pad or the blood continues to be bright red in color, this should be considered serious and you should report to labor and delivery.



Labor Epidurals

LABOR EPIDURALS: What are they and what can I expect? by Brian Kradel, MD

Epidural Analgesia- sometimes called an epidural block causes some loss of feeling in the lower area of your body, yet you will remain awake and alert.

An epidural block is given in the lower back into a small space (epidural space) below the spinal cord. You will be asked to sit or lie on your side with your back curved outward until the procedure is completed. You can move when it is done, but you will not be allowed to get out of bed or walk around.

Before the block is performed, your skin will be cleaned and local anesthesia will be used to numb an area of your lower back. After the epidural needle is placed through this numb area into the epidural space, a small tube (catheter) is inserted through the needle, and the needle removed. Small doses of medication will be given continuously through the catheter to reduce (may not completely eliminate) the discomfort of labor. In some cases, the catheter may touch a nerve during insertion and this may cause a "tingling" or "jerking" sensation down one leg.

Because the medication needs to be absorbed into several nerves, it may take a short while for the epidural to take effect. Pain relief will begin within 10-20 minutes after the medication has been injected.

Although an epidural block will make you more comfortable, you still may be aware of your contractions. You may also feel vaginal exams as labor progresses, but they should not hurt. Your anesthesiologist or nurse anesthetist will adjust the degree of numbness for your comfort and to assist labor and delivery. You might notice temporary heaviness or weakness in your legs. This will disappear after the catheter is removed.

It is important to understand that an epidural may not completely eliminate the discomfort related to labor and delivery. The goal of any epidural placement is to reduce, not eliminate your discomfort.

Although rare, complications or side effects, such as decreased blood pressure, headaches, infection, nerve injury, paralysis and even allergic reaction or death can occur with any procedure. These risks must be accepted and consent signed by you before an epidural is performed.

A woman can decrease some of these risks by holding as still as possible while the procedure is performed. If a headache occurs (risk is 1 in 100), it

often subsides within a few days. If the headache does not stop or it becomes severe, a simple treatment (epidural blood patch) may be needed to help the headache go away. After delivery, your back may be sore from the injection for a few days. However, an epidural should not cause long-term back pain.

The veins located in the epidural space become swollen during pregnancy. Because of this, there is a risk that the anesthetic medication could be injected into one of them. Just like the IV catheter in your hand or arm. If this occurs, you may notice dizziness, rapid heartbeat, a funny taste or numbness around the mouth. If this happens, let your anesthesiologist/nurse anesthetist know right away.

If you require a cesarean delivery, the epidural catheter will be injected with a much stronger drug to increase your pain relief. This will numb your entire abdomen for the surgery. Although there is no pain, there may be feelings of pressure.

The use of an epidural for pain relief during labor and delivery does NOT slow your labor down or increase the chance of you having a cesarean delivery.

No two labors are the same and no two women have the same amount of pain. Some women need little or no pain relief, and others find that pain relief gives them better control over their labor and delivery. Talk to your doctor and your anesthesia expert about your options. Don't be afraid to ask questions and don't be afraid to ask for pain relief if you need it.

Epidurals for labor and delivery are safe procedures if performed by properly trained anesthesia providers (anesthesiologists or nurse anesthetists). Despite the safety of the procedure, it remains an "elective" procedure and not a requirement. It is important that you read and understand the anesthesia consent before the procedure is performed. You must understand and accept the risks.



Hospital Packing List

You may want to pack two bags for the hospital: one for items you'll need during labor and another for everything else you'll need during your stay. Here are some ideas!

FOR LABOR:

- Your insurance card, photo ID, birth plan if you have one, and any hospital paperwork you might need.
- Eyeglasses, if you need them. If you use contacts, you should bring 'backup' glasses just in-case.
- Cell phone, charger, backup battery, tablet, and headphones.
- Diversions for early labor such as a book, magazines, or tablet.
- Robe, nightgown, non-skid socks/slippers, and laboring clothes that you don't mind getting ruined.
- Basic toiletries including hair ties and headbands.
- Pillows for you and for your labor support partner (use colorful pillowcases so yours don't get mixed up with the hospital linen).
- Fragrant oils or lotions for massage. Favorite photos or other focal point. Something to play music on. A hand fan.
- Hard candy, breath mints or strips, lollipops.
- Lip balm or moisturizer.
- A bathing suit for your labor support partner(if you need assistance in the tub or shower.)
- Snacks and a reusable water bottle for you and your labor support person (you may want to pack a small cooler). Change for vending machines.
- Baby book for those tiny footprints!
- Copy of receipt for circumcision if you are having a boy and would like him circumcised.

FOR POSTPARTUM:

- Fresh clothing for you and your labor support person. You may want gowns and sweats. If you are breast feeding, try to find gowns that open or button in the front.
- Cosmetics and perhaps your hair dryer.
- Bring nursing bras and breast pads, lanolin cream and nursing pillow if you plan to breast feed. The hospital will provide sanitary pads and net panties during your stay.
- A going-home outfit. Bring something roomy, comfortable, and easy to get into. Most women still look about 6 months pregnant in that first postpartum week! Plan to wear comfortable flat heeled shoes.

FOR BABY:

- An infant car seat. You can't drive your baby home without one!
- A cute going home outfit (one-piece stretch outfits are easiest).
- A "photo opportunity" outfit if you plan to have any commercial baby photos taken before you go home.
- Comfy newborn clothing if you wish to dress your new baby in anything other than diapers, tee-shirts, and receiving blankets provided by the hospital.
- Receiving blankets 2.
- Mittens, caps, and socks or booties.



- Cash in excess of that needed for snacks
- Work materials
- Pets
- Valuables
- Credit cards or jewelry





Circumcision

If you should have a baby boy and choose to have your son circumcised, our physicians can do this in the hospital, typically 24-48 hours after delivery. At the time of delivery, baby boy is considered an uninsured patient. If you would like baby boy to be circumcised, the uninsured price is \$200 and payment is required before delivery. If the circumcision is covered by your insurance, it is your responsibility to add baby boy to your plan. Once we have been notified by you or your insurance company that he has been added, we will send a circumcision claim to your insurance. Once claim has paid, you will be refunded the entire amount or any difference that may result, dependent on your policy. Circumcision is a minor surgical procedure done to remove the foreskin from the penis. Most baby boys in our culture are circumcised for reasons of personal preference or for religious beliefs. The American Academy of Pediatrics has gone on the record to state that circumcision is not medically necessary and should be performed only for reasons of personal preference. In years past, we believed circumcision was medically necessary to prevent disease. It is now known that this is not the case. However, there is some evidence that HIV and other sexually transmitted diseases are more easily transferred in uncircumcised males. Circumcision carries a small risk of bleeding and infection. There is also a small risk of re-epithelialization in which the foreskin grows back.

WHAT SHOULD I DO RIGHT NOW?

The most important thing to do while you are still expecting is to think it over. Check out a library book, visit web sites for more information, and ask us questions. Try to have your decision made before you deliver. Since many insurers consider circumcision medically unnecessary they do not cover the cost of the procedure. Some Medicaid plans now pay for circumcision. Speak with our Front Desk staff - **now** - for more information regarding your insurance situation, our fees and our payment requirements.

WHAT IF I DECIDE TO HAVE MY BABY CIRCUMCISED?

If we determine that your insurance will not cover circumcision - we ask that you pre-pay our fee before the surgery is performed. We will accept cash, check, money order, MasterCard, VISA or Amex if your insurance doesn't cover it. Our charge for this procedure is \$200. Our providers use the Gomco instrument to perform circumcision. As with any cosmetic surgery, complications can arise that might require additional surgery which may result in additional expense to you. It is best to have circumcisions performed while the baby is in the hospital. If your baby is not circumcised in the hospital, this will need to be performed by a urologist or pediatric surgeon. They will usually not perform the procedure until the infant is 1 year of age.


BASIC CIRCUMCISION CARE

Circumcision care includes being sure to keep your baby's newly circumcised penis clean. If you notice stool on the penis, clean it with warm soapy water. Regular applications of Vaseline will keep healing skin from sticking to the diaper and allows quicker healing. It is important to keep the skin pulled back behind the tip portion of the penis with each diaper change to insure correct healing, especially in the first 3-4 days.

WHAT IF I DECIDE NOT TO HAVE MY BABY CIRCUMCISED?

Your baby will do just fine. You will need to be responsible for teaching your little boy to carefully clean his penis and foreskin every day while bathing. Your pediatrician and the nursery nurses will help you learn how to do this (it is not hard to do). Sometimes little boys become negligent about cleaning (just like trying to avoid brushing their teeth!) and it will be up to you to be sure they develop the habit of cleaning themselves as a part of their daily routine. The risk you face with neglected cleansing of the foreskin is chronic inflammation and a potential for infection. Sometimes this results in a decision to have circumcision done later in life. Late circumcision generally requires hospitalization and can be much more traumatic and painful.

You will need to present the circumcision receipt at the hospital to show it has been paid for.





Postpartum Information

After you leave the hospital, please do not hesitate to call our office if you have questions or problems.

POSTPARTUM OFFICE VISIT

Call the office at (850) 769-0338 to schedule your postpartum appointment. You should be seen in the office in about five to six weeks after your delivery.

You should, however, contact us promptly if any of the following occur:

- Heavy sustained bleeding greater than or equal to one pad per hour.
- Flu-like feeling, fever greater than 100.5, or chills.
- Localized redness or tenderness on your breasts associated with fever.
- Redness, drainage, bleeding at the site of the cesarean incision.
- Foul smelling discharge from the vagina.
- Extreme tenderness of pubic bone, accompanied by frequency, urgency, or burning on urination
- Extreme tenderness in abdomen accompanied by a temperature greater than 100.5.
- Pain that seems to be getting worse, rather than better.

NORMAL CHANGES

In the 4-6 weeks following delivery, the changes of pregnancy are gradually reversed as the body begins to return to its non-pregnant state. The amount of time required for this process varies, depending on the type of delivery you had and other associated medical conditions. The first 6 weeks following the birth of your baby is called the postpartum period.

THE UTERUS

The normal changes of the pregnant uterus to accommodate a growing baby are not reversed overnight. During pregnancy, the uterus increases approximately 11 times its non-pregnant weight, weighing more than 2 pounds immediately after delivery and is about the size of a grapefruit. It can be felt just below the navel. By 8 weeks postpartum the uterus has usually returned to normal size.

AFTERBIRTH PAINS

As the uterus shrinks, its muscle fibers contract, causing afterbirth pains. These contractions are much less painful than labor contractions because there is no associated pain from the cervical dilation. Afterbirth pains are most noticeable the first 3 to 4 days following delivery, particularly for women who have had previous deliveries. These contractions are also pronounced during breast-feeding. Advil, Motrin, or Ibuprofen taken as directed can significantly reduce this pain.

LOCHIA

The drainage from the birth canal following delivery is called lochia. During the immediate few days after the birth, the discharge is like a menstrual flow. In 3-4 days, the discharge becomes more watery and pale. By the second week, lochia is thicker and more yellow in color. Finally, after 4 weeks, the discharge decreases to a minimum as the uterine lining heals. The odor of lochia is usually described a "fleshy, musty, or earthy." The odor should not be bad or offensive. You may have a day of heavier bright red bleeding as the body rids itself of the placenta scar. You may experience occasional cramping and with that, the passing of a clot and brief bleeding. This is normal. Lochia is often heavier when the mother gets out of bed. However, it is important to notify our office if you experience heavy, profuse and persistent bleeding (more than one pad per hour) or if there is a foul odor to the discharge. Please use pads instead of tampons, do not douche, and please abstain from intercourse until the bleeding stops.

VAGINA

The vagina stretches to accommodate delivery and gradually returns to its previous condition by the end of the third week. The supporting structures and muscles of the pelvic floor may not completely return to normal for 6-7 weeks. Episiotomies and lacerations usually require 4 or more weeks to heal. It is important to resume Kegel exercises in the first few postpartum days. These pelvic floor exercises may help speed healing and help muscles return to normal.

It is very common to feel sore and bruised in the vaginal, perineal and rectal areas after delivery. If you have had stitches, these will dissolve gradually and do not need to be removed. It is important to keep the perineal area as clean as possible. After either urination or bowel movement, the area should be rinsed with warm water and gently dried. You may use sitz baths for relief from discomfort (sit in a bath of 6-8 inches of warm water for 15-20 minutes, may repeat 2-3 times a day, may add antibacterial soap to the water). You may use a topical anesthetic spray such as Dermoplast for additional relief. You may take ibuprofen or Tylenol®, two tablets every three to four hours for discomfort.

INCISION CARE FOLLOWING CESAREAN SECTION

Generally, incisions are closed with staples. These are usually removed 5-7 days after the c-section. Call our office shortly after your discharge and set up an appointment for staple removal. The two most important instructions regarding wound care are to keep the incision clean and dry. You should wash the incision daily with soapy water and allow it to completely dry once you are finished bathing. If one or two staples fall out on their own, this is usually not concerning. Sometimes Steri-strips are placed after the staples are removed. It is OK to shower with Steri-strips, if they have not fallen off by 4 days, manually remove them. It is important to notify our office immediately if you notice active bleeding or drainage from the incision. Once again, if the wound is reddening

or if you are experiencing pain not controlled by your pain medications, you need to contact us.

MENSTRUAL CYCLE

The first menstrual period following delivery is usually delayed by breast-feeding. Most bottle feeding patients will experience their first period within 7-9 weeks after delivery. Nursing mothers frequently resume menstrual periods by 12 weeks, however, some do not menstruate until they have completed breast-feeding. Warning: Ovulation will return before the first menstrual period which may result in pregnancy.

INTERCOURSE

It is important to abstain from intercourse until you are cleared by your provider at your postpartum visit. Lubrication may be needed for comfort when you first resume sexual activity. Don't be alarmed if there is discomfort the first few times you have intercourse. Breast-feeding mothers are more likely to have vaginal dryness and some discomfort during intercourse for 4-6 months after delivery. This is caused by diminished estrogen production due to lactation.

CONSTIPATION AND GAS PAIN

If you experience constipation and/or gas pain, try increasing your fluid intake and your intake of fresh fruits and vegetables. Metamucil, Surfak, or Colace may be taken as directed. For gas, you may take Gas-X (simethicone 40 - 80 mg) three to four times daily with meals and at bedtime. Try eating natural foods like prunes, prune juice, apple juice, apple cider, bran flakes or raisin bran. Increase your fluids, especially water. Avoid caffeine (colas, tea, coffee and chocolate). If a laxative is needed, magnesium citrate, Dulcolax, Peri-colace, or milk of magnesia as directed.

HEMORRHOIDS

If you experience hemorrhoids you may relieve discomfort by avoiding standing or sitting for long periods of time, take rest periods whenever possible during the day, lying on your left side with legs elevated on two pillows. Knee/chest position is also pain relieving. Do not allow yourself to become constipated. Drink plenty of liquids, especially water! Sitz baths will help. You may also use Preparation H, Annusol HC, Dermoplast, and Tucks pads (witch hazel), which are available over-the-counter.

DIET

A well-balanced diet is necessary for the healing process. For a breast-feeding mother, the normal caloric requirement is about 2,000 calories per day. Your diet should include foods high in protein and non-saturated fats.

Breast-feeding mothers should increase their fluid intake to 10-12 glasses of water per day to help facilitate adequate milk production. Foods that you eat will end up in the breast milk usually within two to four hours. Keep track of

foods in your diet that may affect either the taste of your breast milk or make your baby colicky. Foods that tend to give you gas will frequently do the same to your baby. We recommend you continue the prenatal vitamins while nursing.

BREAST CARE IN BOTTLE FEEDING MOTHERS

If you are not breast-feeding, we recommend wearing a tight bra and avoiding breast stimulation. Ace wraps and ice packs may also be helpful. You may try applying cabbage leaves to the breast in order to dry the milk supply. Apply the cabbage leaves two to three times daily for about 15 minutes. Call us if you have localized tenderness, redness, or fever.

BREAST CARE FOR BREAST FEEDING MOTHERS

Discuss with your pediatrician issues such as adequacy of the newborn's intake, supplementing with formula, or use of medications while breast-feeding. A firm supportive nursing bra should be worn day and night. Avoid using soap directly on your nipples. Following a breast-feeding session, allow your nipple to air dry. The natural oil glands around the areola will keep the nipples soft and supple. Breast pads should be free of plastic liners to allow air flow. Breast colostrum or milk is an excellent moisturizer for the nipples to prevent cracking and blistering. Breast creams are not advised. Frequent feedings of 8 to 12 times in 24 hours help to prevent engorgement. Nipple soreness is expected for the first one to two weeks. If you experience cracking or bleeding of the nipples, fever, or severe pain, please call the office. Also, consider speaking with a lactations consultant.

***Most insurance plans may cover an electric breast pump under the Affordable Care Act that was passed. Check with your insurance provider about coverage and we can provide you with a prescription if needed.

PLACENTAL INGESTION

Eating the placenta is a trendy practice touted by some who believe there are health benefits for mothers. However, there is no valid research supporting these claims. Most research showing benefit has been funded by the companies that ell placental encapsulation services. The placenta is the temporary organ that grow in the uterus to provide the fetus with nutrients and oxygen. Researchers at Northwestern University School of Medicine reviewed 10 published research studies on the practice. They found no data to support the common claims that eating the placenta either raw, cooked or encapsulated offers protection against postpartum depression, reduces post-delivery pain, boosts energy, helps with lactation, promotes skin elasticity, enhances maternal bonding, or replenishes iron in the body. The researchers also noted that there have been no studies conducted on whether eating the placenta has any potential risks. Emerald Coast OBGYN does not recommend this practice, nor do we offer placenta processing services. The hospital does allow mothers to take their placenta home after the

appropriate consents are signed. If you plan on taking your placenta home, you will need to bring a cooler or waterproof container.

POSTPARTUM BLUES

Regardless of whether you are a first time mother or not, having a newborn is a major life change. Your entire time schedule and priorities will change. In the beginning, at least, your whole world will center on your baby. It is normal on occasion to feel overwhelmed. It sometimes takes 2 to 3 months to establish a routine with your newborn. "Postpartum Blues" are associated with occasional crying, mood swings, and fatigue. These symptoms are very common early on and are usually attributed to sleep deprivation. Don't hesitate asking for help during this time period from friends and family so you can catch up on sleep.

POSTPARTUM DEPRESSION

Less commonly, patients may develop persistent severe emotional symptoms later on in the post partum period. Post partum depression is characterized by:

- Excessive crying
- Inability to sleep or excessive sleeping
- Loss of appetite
- Fear and anxiety
- Feelings of hopelessness or loss of control
- Hostility or self-blame
- Difficulty concentrating or making decisions

These symptoms are not signs of weakness or inadequacy. You should seek professional help (our office or a therapist) if you are experiencing these symptoms. Treatment may include medication, counseling, and in severe cases hospitalization. With proper treatment, most women recover fully. Above all, remember that postpartum depression is a real condition and help is available.

NARCOTIC MEDICATION DISPOSAL

DEA regulations do not permit practitioners to dispose of abandoned controlled substances, even if the ultimate user is a patient of the practitioner. Please note. Do NOT accept unused controlled substances from your patient, even if you were the prescribing provider. Advise the patient to return the unused medications to their pharmacy. We are NOT authorized to receive them. We have disposed of patients unused controlled substances in the past. However, DEA regulations now prohibit this.

Family Planning



In today's world we are very fortunate to have many contraceptive options available to us! Perhaps never before have so many choices been within our reach. We divide them into temporary and permanent. The temporary methods are further divided into hormonal, mechanical, and barrier. Be sure to talk with us about your plans. We can help you sort out the best option for you.

PERMANENT METHODS

For permanent contraception we are generally speaking about sterilization procedures such as tubal ligation for women and vasectomy for men. **Remember, no sterilization procedure has a 0 % failure rate.** The failure rate generally quoted to patients undergoing tubal ligation is about 1 in 100. The failure rate is age dependent with younger patients having a higher rate of unintended pregnancy after sterilization.

TUBAL STERILIZATION

Tubal sterlization procedures are done surgically by interrupting the tubes between the ovary and the uterus. This interruption prevents the meeting of the woman's egg and the man's sperm. Tubal sterilization will make you sterile and should never be seen as a temporary or easily reversed procedure. At Emerald Coast the primary method used is placement of small metal surgical clamps placed on the fallopian tubes to seal them. Tubal sterilization is always a surgical procedure and must be done at the hospital or at one of the outpatient surgical centers. In patients with a family history of ovarian cancer, it may be preferable to remove the entire tube. The procedure is done on an outpatient basis unless you have it scheduled as part of a planned Cesarean section or as a postpartum procedure done shortly after delivery. If not done at the time of c-section, patients will usually undergo general anesthesia (you are put to sleep). Postpartum tubal sterilization is done through an incision under your navel. The incision is slightly larger for an immediate postpartum tubal sterilization, so if the cosmetic appearance of your belly button is important to you, a laparoscopic procedure done 8 weeks postpartum will probably be a better choice for you.

Some insurance providers, such as Medicaid, require that initial paperwork be started at least 30 days before you plan to have the tubal sterilization performed. For this reason, we strongly recommend that you initiate the paperwork at about 28 - 32 pregnancy weeks if you believe you are seeking a permanent procedure. Be sure to let us know so that we can help you make plans. Remember, you can begin the required paperwork and still change your mind at any time before the tubal sterilization is done.

VASECTOMY

Vasectomy is a minor but permanent outpatient office procedure for men and is available through a urologist. If your partner is considering a vasectomy, please ask us for referral information.

TEMPORARY METHODS

Temporary methods: Of course, any of the temporary methods will provide contraception but only if used correctly and consistently!

HORMONAL

- Oral contraceptive pills
- Depo Provera injections given every 3 months
- The vaginal ring inserted monthly
- The patch applied weekly to skin.
- The hormonal intrauterine device good for up to 5, 8 or 10 years (depending on the device)
- The single rod implant/Nexplanon good for up to 3 years.

We can provide you with information about each of these methods and discuss with you which method may suit your individual situation best. These methods are all very effective - when used as intended. Each option carries its own risks. We are happy to discuss the advantages, risks, and limitations of any of these methods with you.

NON-HORMONAL

These will generally include barrier methods such as diaphragms and condoms, spermicidals, the non-hormonal intrauterine device, and natural family planning methods such as ovulation charting. Be aware that breast-feeding and penis withdrawal will provide uncertain results and cannot always be counted on to prevent pregnancy. We are happy to discuss the advantages and limitations of any of these methods.

INTRAUTERINE DEVICES (IUD's)

These are small "T" shaped devices that are placed into the uterus by your provider in the office. The procedure is usually quick and associated with only mild cramping. There are several now available. Two types are:

- **Paragard:** This IUD has no hormones. The device has a copper coating and primarily works by incapacitating sperm and may be used for up to 10 years.
- Mirena/Liletta: These IUD's have a progestin (hormonal) coating that works by disrupting sperm, thinning your uterine lining, and thickening cervical mucus. One beneficial side effect of this device is lighter periods. It can take 4-6 months until patients see a decrease in their periods, and often during this time they are irregular. These devices may be used for up to 8 years.

- **Kyleena:** This IUD is good for up to 5 years and is an estrogen free device. It also works by thickening cervical mucus, inhibiting sperm movement, reducing sperm survival, and thinning the lining of your uterus. It is a smaller device and may be preferable for patients who had a c-section.
- Emergency Contraception: Any of us can run into this type of emergency. What if the condom breaks, or you have not consistently taken your contraceptive? What if you suddenly remember the antibiotics your dentist prescribed? The Plan B emergency contraceptive is now available over-the-counter (without prescription) for all women. Emergency contraception is not as certain as the non-emergency or continuous methods of birth control. However, when taken properly within 72 hours of unprotected sex, Plan B emergency contraception will be effective about 80% of the time. Plan to spend \$40 to \$45 for Plan B. Most pharmacies carry this product over the counter.



Interesting Websites

INTERESTING WEBSITES YOU MAY WANT TO VISIT**

www.emeraldcoastobgyn.com

Our website will have clickable links if you want to use it as a starting point.

PREGNANCY

www.4women.gov/pregnancy www.marchofdimes.com www.childbirthconnection.org www.cdc.gov www.lamaze.org www.acog.org/publications/patient_education www.mymidwife.org www.osteopathic.org www.spinningbabies.com

PLACENTA DONATION

www.donateyourplacenta.org

CIRCUMCISION

www.nlm.nih.gov/medlineplus/circumcision.html www.mayoclinic.com/health/circumcision/PR00040

BREASTFEEDING

www.lalecheleague.org www.kellymom.com

BIRTH CONTROL

 $www.KnowMyCycle.com\\www.gynob.emory.edu/familyplanning/contraception.cfm$

FDA website:

www.fda.gov/womens/healthinformation/birthcontrolguide.pdf

** Emerald Coast OB/GYN is not responsible for content or advice found on any internet website other than our own. If you have concerns or questions about advice found on an internet website, please discuss it with us.


Pregnancy Myths



- 1). **Have a baby lose a tooth.** Not true. A pregnant woman should never lose a tooth due to her baby! Preventive and emergency dental care is important during pregnancy. Diligent dental care is important in pregnancy. You should strive to have 1800mg calcium from foods or supplements daily during pregnancy.
- 2). Taking iron while pregnant will cause your baby to have dark skin. Not true. Baby's skin color is caused by family genes, not what you eat.
- 3). Sex during pregnancy will harm your baby. Not true. Sex is safe during all trimesters unless otherwise specified by your care provider.
- 4.) Raising your arms above your head will cause your baby to become tangled in the umbilical cord. Not true. All babies swim, twist, and turn within their sac of amniotic fluid. A mother's activity will not cause an umbilical cord problem. Approximately 25% of all healthy babies will be born with their umbilical cord wrapped about their shoulders or necks. Sometimes the cord may even form a knot. There is a gel like substance in the cord that protects it from compression.
- 5.) Having your hair dyed or permed during pregnancy will harm your baby. Not true. Chemicals used in today's hair products are generally not harmful to your baby.
- 6.) Eating for two means eating twice as much. Not true. While you and your baby do need a healthy diet this does not mean eating twice as many calories! You actually should only eat a total of 300 calories/day more during your pregnancy.
- 7.) Painting the nursery can harm your baby. Maybe or maybe not... It is certain that scraping old lead based paint off your woodwork can cause harm so let others do the prep work if the old paint was applied before 1977 (when lead paints were taken off the market.) When painting, be sure to have good ventilation and use fans. Avoid oil-based paints and products containing mercury. Avoid solvents. Use latex paints that do not contain ethylene glycol ethers or biocides. Better yet get others to do the painting for you!
- 8.) If you have lots of heartburn, your baby will have a full head of hair. Not true. There is no evidence that heartburn in the mother is related to hair growth in the baby.
- 9.) Sex determination myths. All false!!! We cannot determine gender by fetal heart rate, the Drano test, the pencil test, maternal acne, carrying high, carrying wide, carrying low, or severity of nausea & vomiting.
- 10.) **Immersion in water will harm my baby.** Not true. Warm baths are soothing and can help relieve muscle aches and pains. Bath water will not enter your vagina and cause infection in your baby. It is best to avoid overheating by

keeping bath temperature no higher than 100 degrees. Also, it is best to avoid bathing when at home alone during the later weeks of your pregnancy so that someone can help you out of the tub to reduce the chance of a slip or fall.

- 11.) You will harm your baby if you sleep on your back. Relax! It is true that you will have improved circulation to your kidneys and uterus if you avoid sleeping on your back after the 20th week of pregnancy. When your uterus becomes large it can squeeze the blood flow trying to return to your heart and this can also cause you to feel dizzy or nauseated. After your 20th week try to rest and sleep on your side. If you wake up and find yourself on your back, don't worry just reposition yourself to the side.
- 12.) Touching kitty litter will harm your baby. Not true. It is not the litter we worry about. It is the feces of outdoor cats that eat birds. Their feces can contain toxoplasmosis which is a rare but dangerous parasite. Indoor cats that eat packaged cat food will rarely have this organism. There are several solutions to this concern. Your vet can check your pet for toxoplasmosis to see if there is any threat to you and your baby or simply have someone else change the litter box. If the pregnant woman must clean the litter box she should wear gloves and a mask. As always, hand washing is important when handling any pets or after outdoor gardening work.
- 13.) You are more likely to go into labor on the night of a full moon or during a storm. Labor is triggered by a complex series of hormonal/biochemical signals that have no known connection to moon phases, atmospheric conditions, or the position of the stars.
- 14.) Water birth will make my baby's birth transition easier. Once the baby is out, the placenta starts to separate which means less oxygen through the umbilical cord. Babies need air! This being said, soaking in water is usually very comforting during labor. Just be sure to get out of the bathtub before you deliver.



Cord Blood Banking



Cord blood refers to the few ounces of blood remaining in the umbilical cord and placenta which has traditionally been discarded as medical waste following the birth of a child. Physicians and scientists discovered that cord blood contains stem cells (similar to that in bone marrow) that have the ability to replicate or develop into additional cells which can be used to treat life-threatening diseases.

Cord blood banking refers to storing the cord blood which remains in the umbilical cord and placenta after your baby is delivered and safely in your or the pediatric teams' care. The collection of cord blood does not harm the mother or the baby.

The cord blood banking process consists of collection, shipment to the lab, processing, and cryopreservation of the cord blood. Cord blood transplantation is the process where the stored cord blood is thawed and subsequently infused in patients for treatment of more than 70 life threatening diseases.

Additionally, there are many promising clinical trials being performed using cord blood to treat diseases and conditions such as diabetes (types 1 & 2), cerebral palsy, cartilage defects and spinal cord injuries, to name a few. The existence of these trials does not guarantee that they will become clinical therapies. Currently, the chances of a child or family member being treated with a unit of cord blood are about 1 in 2,700. This number is expected to change as technology for its use continues to improve. Parents should also be aware that if a child were to develop leukemia that cord blood would not be able to be used since the stem cells will have the same genetic markers of leukemia.

Patients considering private cord blood banking should be informed of the following issues:

- The indications for cord blood transplant are limited to certain genetic, hematologic, and malignant disorders. For a complete list go to <u>www.nmdp.org</u>
- 2. Routine storage of umbilical cord blood as "biologic insurance" against future diseases is not recommended by the *American Congress of Obstetricians and Gynecologists* nor the *American Academy of Pediatrics*.

- 3. Cord blood banked by families is much more likely to be used by the donors' siblings than the donors.
- 4. Private storage of umbilical cord blood should be considered if there is a family member with a current or probable future need for transplantation.
- 5. Cord blood stored in a private cord blood bank is saved for that family. Cord blood has approximately a 30% probability of being an exact match or very close match for each brother or sister. A family that chooses to store their baby's cord blood pays a fee for the collection, cryopreservation, and annual storage of the umbilical cord blood. Selection of a private cord blood bank is an important decision. Parents should read all contracts thoroughly and inquire about quality measures. *Florida Woman Care* has selected *Cord Use* as the preferred cord banking company based on the quality of the transportation, cryopreservation techniques, and customer service. www.corduse.com
 - 6. Parents do have the option to donate cord blood to a public cord blood bank. Cord blood donated to a public cord blood bank is available for any patients who need a transplant. The donation process is free to the parents donating the umbilical cord blood. Today, however, only certain hospitals are able to collect umbilical cord blood for possible storage in public cord blood banks. Cord blood donated to public banks but does not meet criteria for the public inventory can be used for ethical research to improve transplantation or develop new therapies using cord blood.

PLACENTA DONATION

Gulf Coast Regional Medical Center offers a program where you can donate your placenta through Regenerative Biologics. If you are having a c-section, ask your provider about this program. By donating your placenta, you can help up to 100 eyes see again that were damaged by disease or accident. Also, The natural healing properties from your placenta have a multitude of surgical benefit. There are no side effects or risks associated with the donation of your placenta to either your self or your newborn, and there is no cost involved. To learn more, visit their website at www.donateyourplacenta.org.

Notes





EMERALD COAST OB/GYN

wishes to thank **Color Press Printing** for their outstanding efforts in the development of our OB Booklet. We could not have completed the book without their creative expertise and commitment to our project.



www.colorpressprinting.com

3430 Highway 77 Suite D • Panama City, FL 32405

P: 850.763.9884 • F: 850.769.6999

We at Emerald Coast OB/GYN would like to thank those who have taken out advertisements in support of our book and patient education. We are truly appreciative. Emerald Coast OB/GYN has no financial association with any of our advertisers.



EMERALD COAST CLINICAL RESEARCH LLC

Clinical Research Studies

One of the benefits of being an Emerald Coast OBGYN patient, is the opportunity to participate in important clinical trials advancing women's health. By being part of a clinical trial, participants may find out about new treatments before they are widely available. Some studies are designed for, or include, people who are healthy but want to help find ways to prevent a disease, such as one that may be common in their family. The sponsors of clinical trials often compensate participants financially.

Please talk to your ECOBGYN provider about our current studies that might interest you!



Panama City Surgery Center offers you the top medical specialties & services in one convenient location. Outpatient — also known as ambulatory surgery — does not require an overnight hospital stay.

Our Specialties Include:

Gastroenterology – General Surgery - Laser Surgery - OB/GYN - Oral & Maxillofacial Surgery Orthopedic Surgery - Otolaryngology - Pain Management - Plastic Surgery – Podiatry – Robotic Surgery Urology

We are licensed by the state of Florida and approved by the Centers for Medicare and Medicaid Services. Panama City Surgery Center is inspected and accredited by The Joint Commission, which is a National Accreditation Agency.

Address: 1800 Jenks Avenue, Panama City, Florida 32405 | Phone: (850) 769-3191

DR. AMY BELL DOCTOR OF CHIROPRACTIC

MOTHER OF TWO YOUNG CHILDREN & PASSIONATE ABOUT DELIVERING TOP NOTCH CARE

- SPECIALIZES IN PRENATAL, POSTNATAL & PEDIATRIC CARE
- UTILIZES SAFE, GENTLE AND EFFECTIVE ADJUSTMENT TECHNIQUES
 - CERTIFIED IN THE WEBSTER TECHNIQUE

BENEFITS OF CHIROPRACTIC CARE DURING PREGNANCY

- PROPER PELVIC BALANCE AND ALIGNMENT FOR YOU, PROPER POSITIONING FOR BABY
 - IMPROVED NERVOUS SYSTEM FUNCTION
 - FASTER LABOR AND DELIVERY
 - RELIEF OF LOWER BACK PAIN
 - IMPROVED RECOVERY



850-704-0008

2318 ST ANDREWS BLVD PANAMA CITY, FLORIDA 32405 T: 850-704-0008 F: 850-250-2582 DRAMY@ELITECHIROPC.COM



Are you or someone you know pregnant or recently had a baby?





Now more than ever it is crucial that pregnant women and new parents are offered the support and assistance they need to navigate their pregnancy. CONNECT helps link pregnant women and infants to the home visiting service that suits their needs and will benefit their families the most. FREE support will be provided over the phone to keep you and your family safe.

Services Include:

- Pregnancy, Childhood, and Parenting Education
- Breastfeeding Education and Support
- Help to Quit Smoking
- Infant Milestones and Child Development
- Family Planning
- Counseling
- Financial Planning and Links to Community Assistance
- Referrals to Other Community Resources
- Infant and Child Safety (Car Seat Safety, Safe Sleep, and More!)

Home Visiting Programs in Bay, Franklin, & Gulf Counties:







To refer yourself or make a community referral for someone you know, call:
For Bay County Residents; 850-527-3745
For Gulf & Franklin County Residents: 850-370-0006





The Vagicians are here for the typical female, answering not-so-typical questions about women's health and well-being.

SEASON 1 immediately available on Sound Cloud, iTunes, Spotify, & YouTube!

Please tune in on any of these outlets for the latest episodes.

Dr. Wolf says, "Have a Vagical Day!"

Panel of experts include:

Dr. Sam Wolf, Dr. Jacob Martin, & Dr. Roslyn Mallory





Microblading





The newest technique to achieve gorgeous eyebrows! It is the semi-permanent solution to creating the perfect brows. Also known as "3-D brow tattooing", it is so much more than just giving a tattoo – it is the art of using a special blade to create intricate, hair-like strokes to give the illusion of having beautifully full, natural looking eyebrows.

Beautiful eyebrows are something nearly every woman wants.

The perfect brows frame the face and bring out the eyes.









We offer eyebrow Microblading and Shading, Permanent Lip blushing (tattoo), Eyeliner, 3D areola/nipple tattoo for breast cancer survivors and Scalp Micropigmentation (hairline tattoo).

All of our artists are highly trained in the most advanced permanent makeup techniques, licensed by the Department of Health in Bay County, and fully-insured.

Microblading in Panama City is now available for you!

Quit wasting precious time drawing on your eyebrows and spending endless amounts of money on expensive eyebrow gel/pencils by simply calling our Microblading Specialists at Brow Artistry by Irina.









Text or call (850) 774-4418 to schedule your appointment today!



Family Practice (6 months and up)

Well Checks

IV Hydration

Physicals

Wellness





WWW.FAITHFAMILYWELLNESS.HEALTH

850.788.3120

#WOMENOFWELLNESS #WHOLEFAMILYHEALTH





Helping You Live healthier

Experience the

Rx Express Difference

Your hometown pharmacy with a personal touch



Lauren Parker PharmD, Staff Pharmacist

RETAIL AND COMPOUNDING PHARMACY

"Your Vaccine Destination"

Offering Covid, Flu, DPT, Shingles, and Hepatitis Vaccines

rxexpresspc.com reexpresspc@gmail.com

850.215.7455

3104 W 23rd Street - Panama City FL 32405

BC BS of Florida Medicare D Rx Pharmacy Express of Panama City

PROUDLY SUPPORTS OUI





If you're having a C-section, ask your doctor about our Placenta Donation Program

By donating your placenta, you can help up to 100 eyes see again that were damaged by disease or accident.

Donated tissue is used in nerve repair, wound care, orthopedic recoveries, and head trauma cases.



- No cost to patient
- No risk to mother and baby
- Doesn't interfere with cord blood donation/banking

"I felt honored and proud to help. It was an amazing experience for me and will be a great story to tell my baby about"

Donor mom, Daisy C., Gainesville, FL

Learn more about donation at www.donateyourplacenta.org or contact us:

Phone: 866.432.1164





OUR SERVICES:

Obstetrical Services 3D Mammography

Davinci Surgery System

We feature the Hologic® Genius™ 3D MAMMOGRAPHY machine. This technology delivers a series of detailed breast images, allowing your doctor to better evaluate your breasts layer by layer.

Through the use of the da Vinci® SURGICAL SYSTEM, surgeons are now able to offer a minimally invasive option for complex surgical procedures including hysterectomies that might otherwise require an open abdominal incision.

103 E. 23rd Street • Panama City, FL • 850.769.0338 emeraldcoastobgyn.com

BALDWIN pediatrics

Baldwin Pediatrics offers:

- Free Prenatal Consultations
- Newborn Care
- Hospital Nursery Rounds
- Well Baby Visits
- Immunizations
- Same Day Sick Appointments
- Saturday/Extended Hours
- Asthma/Allergy Treatment



Our Providers



Dr. Rubina Azam MD, FAAP, ABIHM

- Dr. Roland Jaoude
- Summer Moran, NP
- Jennifer Slaton, NP
- McKenna Devlin, PA

Call today to schedule a consultation!

850-522-1522




528 W. Baldwin Road, Panama City, FL 32405











- Free Prenatal Consulting
- Newborn & Premature Baby Check-up
- Lactation Consultation Available
- Vaccinations
- All Newborn, Childhood,
 & Adolescent Illnesses
- Well-Child Check-up
- School Physicals
- Asthma & Allergy
- ADHD
- Establishing Pediatric Care





Panama City Office:

621 W Baldwin Road, Suite A Panama City, FL 32405 (850) 747-3661

Panama City Beach Office:

12907 Panama City Beach Pkwy, Suite 201 Panama City Beach, FL 32407 (850) 238-8720

Monday-Thursday 8am-5pm Friday 8am-4pm

After Hours Clinic (Urgent Care)

Monday-Thursday 5pm-8pm Friday 4pm-7pm Saturday 9am-2pm

Moms LUVUs

Hear what moms have to say about us

"They knew exactly the support we needed in those hours — and they provided it with expertise and compassion."



